

# Skeletal Muscle Fatigue: Cellular Mechanisms

D. G. ALLEN, G. D. LAMB, AND H. WESTERBLAD

*School of Medical Sciences and Bosch Institute, University of Sydney, Sydney, New South Wales, and Department of Zoology, La Trobe University, Melbourne, Victoria, Australia; Department of Physiology and Pharmacology, Karolinska Institutet, Stockholm, Sweden*

---

|   |     |
|---|-----|
| I. Introduction   | 288 |
| A. Continuous maximal activity  | 289 |
| B. Repeated short tetani  | 289 |
| C. Delayed recovery from fatigue  | 290 |
| II. Sequence of Excitation-Contraction Coupling in Muscle                                   | 290 |
| III. Metabolic Changes in Working Muscles   | 291 |
| IV. Differences Between Fast and Slow Fiber Types   | 292 |
| V. How to Study Fatigue?  | 292 |
| A. Models of fatigue and their limitation   | 292 |
| B. Muscle temperature   | 294 |
| VI. Excitability and Extracellular K <sup>+</sup> Accumulation                              | 295 |
| A. Basis of excitability  | 295 |
| B. Alterations in excitability  | 296 |
| C. Factors helping to prevent loss of excitation during normal exercise                     | 297 |
| VII. Metabolic Changes and Fatigue  | 299 |
| A. Inorganic phosphate  | 300 |
| B. Lactate and H <sup>+</sup>   | 303 |
| C. ATP and Mg <sup>2+</sup>   | 305 |
| D. Glycogen   | 306 |
| VIII. Reactive Oxygen Species   | 307 |
| A. ROS production is accelerated in active muscles  | 308 |
| B. Sources of ROS in muscle   | 308 |
| C. ROS scavengers reduce fatigue  | 309 |
| D. Mechanisms by which ROS cause fatigue  | 309 |
| IX. Ca <sup>2+</sup> Compartments and Movements: Role in Fatigue                            | 311 |
| A. SR Ca <sup>2+</sup> content and Ca <sup>2+</sup> release in fast- and slow-twitch fibers | 311 |
| B. Effects of SR content changes on Ca <sup>2+</sup> release                                | 311 |
| C. Ca <sup>2+</sup> content changes with repeated stimulation and with exercise             | 312 |
| D. Mitochondrial Ca <sup>2+</sup> handling  | 312 |
| X. Shortening Velocity and Mechanical Power   | 313 |
| XI. Slowing of Relaxation   | 314 |
| XII. Factors Counteracting Fatigue  | 315 |
| XIII. Recovery From Fatigue   | 315 |
| A. Delayed recovery from fatigue  | 316 |
| B. Prolonged changes in the contractile apparatus   | 317 |
| C. Prolonged reduction in Ca <sup>2+</sup> release  | 318 |
| XIV. Blood Flow and Intracellular Po <sub>2</sub>   | 318 |
| XV. Concluding Remarks  | 319 |

---

**Allen DG, Lamb GD, Westerblad H.** Skeletal Muscle Fatigue: Cellular Mechanisms. *Physiol Rev* 88: 287–332, 2008; doi:10.1152/physrev.00015.2007.—Repeated, intense use of muscles leads to a decline in performance known as muscle fatigue. Many muscle properties change during fatigue including the action potential, extracellular and intracellular ions, and many intracellular metabolites. A range of mechanisms have been identified that contribute to the decline of performance. The traditional explanation, accumulation of intracellular lactate and hydrogen ions causing impaired function of the contractile proteins, is probably of limited importance in mammals. Alternative explanations that will be considered are the effects of ionic changes on the action potential, failure of SR Ca<sup>2+</sup> release by various mechanisms, and the effects of reactive oxygen species. Many different activities lead to fatigue,

and an important challenge is to identify the various mechanisms that contribute under different circumstances. Most of the mechanistic studies of fatigue are on isolated animal tissues, and another major challenge is to use the knowledge generated in these studies to identify the mechanisms of fatigue in intact animals and particularly in human diseases.

## I. INTRODUCTION

Muscles that are used intensively show a progressive decline of performance which largely recovers after a period of rest. This reversible phenomenon is denoted muscle fatigue. The phenomenon must have been recognized by perceptive observers since posterity, but studies of the mechanism are relatively recent. Needham (1971) (328) provides a comprehensive account of the history of muscle contraction and metabolism and quotes Berzelius (1807) for the discovery that the muscles of an exhausted stag contained lactic acid. Mosso (1904) (318) introduced the experimental study of the topic with figures illustrating the rapid fatigue in humans when a finger lifts a heavy load. He showed that the rapid fatigue occurs even when the nerve was stimulated electrically, implying that fatigue is in the muscle rather than in the central nervous system. A seminal contribution was made by Hill and Kupalov (1929) (208) who showed that an isolated frog muscle stimulated in  $N_2$  gas fatigued rapidly and accumulated lactic acid. A striking result was that if the muscle was then transferred to a  $N_2$ -saturated Ringer solution, the performance recovered as lactic acid diffused out of the muscle, suggesting that lactic acid could cause fatigue. Eberstein and Sandow (1963) (149) were the first to suggest that failure of excitation-contraction (EC) coupling contributed to muscle fatigue by showing that a fatigued muscle could recover much of its force when perfused with caffeine, known to directly facilitate re-

lease of  $Ca^{2+}$  from the sarcoplasmic reticulum (SR). Burke et al. (1973) (75) stimulated individual motor units in cat muscles to exhaustion and identified the muscle fibers involved by the depletion of glycogen. These influential studies showed that fast fibers fatigued extremely quickly, whereas slow fibers were essentially unfatigable. The development of the muscle biopsy technique by Bergström et al. (1967) (46) and the application of muscle NMR to fatigue by Dawson et al. (1978) (117) accelerated the understanding of the biochemical changes during fatigue.

One definition of fatigue is any decline in muscle performance associated with muscle activity. This is particularly clear when maximum isometric force is measured in repeated tetani (Fig. 1) and shows that there is a progressive decline that is visible even on the second tetanus of the series. Other aspects of muscle performance also change during fatigue, notably shortening velocity is reduced and the time course of relaxation slows. Most practical activities are dependent on the power output of the muscles involved and, since power output is the product of both force and shortening velocity, the decline in performance can be larger than the decrease in isometric force. Of course, the decline in performance is not immediately apparent if a submaximal activity is performed, and in this situation fatigue manifests itself eventually as a failure to be able to continue the activity at the original intensity (48), often called exhaustion. In such an activity, the progress of fatigue can




FIG. 1. Force records during fatigue produced by repeated short tetani in an isolated mouse flexor digitorum brevis (FDB) fiber; each tetanus appears as a vertical line. In the *top panel*, the phases of fatigue (see sect. vii) have been indicated. The *bottom panel* shows records from the same fiber fatigued in the presence of cyanide to inhibit mitochondrial oxidative phosphorylation. Stimulation protocol: 350-ms, 70-Hz tetani repeated every 4 s for 2 min, and the interval was decreased by  $\sim 20\%$  every 2 min (interval changes indicated by open triangles). Temperature was  $25^\circ\text{C}$ . [From Lännergren and Westerblad (268).]

be estimated by occasionally interpolating a maximal contraction.

In voluntary contractions, muscles are activated by complex pathways starting in the cortex and leading to excitation of lower motor neurons in the spinal cord. The axon of the lower motor neuron carries the action potentials to the neuromuscular junction of the muscle. For simplicity, the processes inside the spinal cord and above are defined as central, whereas the processes in the peripheral nerve, neuromuscular junction, and muscle are defined as peripheral. Clearly fatigue can potentially arise at many points in this pathway and can usefully be divided into central and peripheral fatigue. Early studies by Merton (1954) (308) suggested that in well-motivated individuals, the fatigue in a small muscle of the hand could be entirely peripheral. Later studies suggest that a small degree of central failure of activation often occurs during maximal activation of muscles and that during fatigue there is often a substantial central component (for review, see Gandevia, Ref. 179). Nevertheless, it is universally agreed that much of fatigue arises in the muscles and can therefore be studied in isolated muscle tissues. This review focuses on the components of fatigue that lie within the muscle.

It is usual to make a distinction between muscle fatigue and muscle injury, although undoubtedly the two phenomena overlap. Fatigue is usually defined as the reversible decline of performance during activity, and most recovery occurs within the first hour. However, there is also a slowly reversible component that can take several days to reverse (155). Muscle injury also causes a decline in performance that reverses only very slowly. Muscles that are stretched during contraction (eccentric contractions) are particularly prone to injury or damage. Injury is characterized by structural abnormalities including sarcomeric disorder, membrane damage resulting in the loss of soluble enzymes such as creatine kinase, and inflammatory processes including cytokine release and phagocytic cell infiltration. Recovery from the most serious injuries involves activation of satellite cells and regeneration of damaged fibers. Muscle damage has been reviewed recently (12, 363) and will not be considered here.

### A. Continuous Maximal Activity

If a muscle is stimulated continuously at a frequency close to that which gives maximal force, then force production generally shows a rapid decline (Fig. 2) often called high-frequency fatigue (52, 227). A characteristic of this type of fatigue is that the recovery is also very rapid, often having a component of recovery with a time course of only 1–2 s (52, 227). In a human, this type of contraction would be encountered in lifting a very heavy object




FIG. 2. Force record from a fast-twitch lumbrical muscle fiber of *Xenopus* stimulated at 70 Hz for 30 s. Recovery followed with 500-ms, 70-Hz tetani given at 2-s intervals. Note very rapid recovery after 2 s. [From Lännergren and Westerblad (265).]

(e.g., a piano), and it is generally recognized that onset and recovery of fatigue are rapid in this situation. In intact individuals, an important consideration is that when force exceeds ~50% of maximum, then the muscle circulation collapses (33). Consequently, continuous maximal contractions occur in a nonperfused (ischemic) muscle, though perfusion would be present during recovery. Studies of the firing rate of motor neurons in intact individuals show that in such near-maximal contractions, the initial firing rate is high but that this rate falls steadily over a minute or so (50). In experimental studies of high-frequency fatigue, it is common to simply stimulate at a constant high frequency, so it is important to realize that such studies do not accurately reproduce what happens in the intact individual.

### B. Repeated Short Tetani

A popular stimulation pattern for the study of fatigue is repeated, short tetani (Fig. 1). Unfortunately, there is little uniformity in the detailed protocols used by different groups. One key variable is the fraction of the time during which the muscle contracts (duty cycle), which is commonly between 0.1 and 0.5. Another key variable is the stimulus frequency during the tetani which determines the degree of activation and will influence the rate of decline of force. This pattern of stimulation, which obviously simulates many natural activities (e.g., walking, running, and breathing), leads to a much slower rate of fatigue than the continuous high-frequency stimulation described above. Commonly the fatigue protocol is stopped after a fixed number of tetani or when the force reaches a certain predetermined level such as 50% of the initial tetanic force. The rate of recovery from this pattern of activity is quite variable. There is usually a phase of recovery that is nearly complete after 5–10 min and sometimes a much slower component (discussed below). Usually recovery of maximum force is virtually complete by 30 min, and it is possible to perform repeated fatigue runs

with nearly similar time courses after a recovery period of >30 min. This allows the comparison of fatigue runs under different conditions within the same fiber avoiding interfiber variability (90, 203).

### C. Delayed Recovery From Fatigue

As noted above, after repeated short tetani there may be a very slow component of recovery. This phenomenon was first described by Edwards et al. (155), who used repeated voluntary contractions in humans under ischemic conditions continued until the force was negligible. Recovery measured in brief tetani at high frequencies (50–100 Hz) was relatively fast ( $t_{1/2} \sim 5$  min), whereas recovery at low frequencies (10–20 Hz) was very slow ( $t_{1/2}$  1–2 h) with a small component of weakness still persisting after 1 day. They called this phenomenon “low-frequency fatigue,” a name we do not recommend for the reasons discussed below. The phenomenon is particularly prominent if the exercise contains a component of muscle stretch (114), which is liable to cause muscle damage, and this is one area in which muscle damage and fatigue appear to overlap. For human subjects, the force is reduced in a frequency range that corresponds to the firing frequencies of human motor units during low- to moderate-force voluntary movements (300), and the phenomenon is probably responsible for the feeling of weakness that can persist for several days after a period of intense exercise. Presumably to achieve the necessary force for a particular activity, the brain must increase the firing rate or activate more motor units for a particular muscle and interprets this information as weakness, even though the muscle may show no decrement in maximal force.

As noted above, high-frequency fatigue is widely used to describe fatigue resulting from continuous maximal contractions. This suggests by analogy that “low-frequency fatigue” should be used for continuous stimulation at low frequencies, but in fact, Edwards et al. (155) used the term to describe a type of recovery that was slower when measured at low frequencies. Thus we recommend using the term *prolonged low-frequency force depression* for this phenomenon and avoid the term *low-frequency fatigue* that is now used for several quite different situations. Furthermore, high-frequency fatigue is sometimes used to describe fatigue induced by repeated short tetani, where the stimulation frequency is high during the tetani. Thus this term may also be misinterpreted, and we recommend that it is also avoided.

## II. SEQUENCE OF EXCITATION-CONTRACTION COUPLING IN MUSCLE

The chain of events involved in EC coupling in skeletal muscle is now reasonably well understood (for re-

view, see Refs. 140, 306, 424). Obviously a failure anywhere in this chain of events could contribute to fatigue. The muscle action potential (AP) is initiated at the neuromuscular junction by release of acetylcholine as a consequence of the AP in the motor neuron. The neuromuscular junction appears to provide uniform 1:1 AP transmission under physiological conditions (179), and failure at this point will not be considered in this review. Transmission of the AP along the surface membrane of the muscle fiber depends on local currents activating distant  $\text{Na}^+$  channels. Potentially AP transmission is influenced by numerous factors including the membrane potential, the  $\text{Na}^+$  and  $\text{K}^+$  concentrations extracellularly and intracellularly, the internal and external resistances, and the membrane resistance and capacitance. Many of these factors can vary in fatigue, and possible contributions of the AP transmission to fatigue are considered in section vi. The AP is actively conducted down the transverse tubules (t tubules) into the interior of the muscles (47). Because of the small volume of the t-tubular system, changes in ionic concentrations are particularly prominent in the t tubules. However, because there is no simple way to measure the AP within the t-tubular network, assessing the role of the T system is difficult. The t-tubular membrane expresses high levels of L-type  $\text{Ca}^{2+}$  channels (or dihydropyridine receptors, DHPRs, or voltage sensors) which change their conformation during an AP, resulting in charge movement (397). The voltage sensors are in close contact with the SR  $\text{Ca}^{2+}$  release channels (the ryanodine receptors, RyR), which in mammalian skeletal muscle are principally the RyR1 isoforms. Charge movement in the voltage sensors results in opening of the RyR receptor and SR  $\text{Ca}^{2+}$  release. The mechanism of this interaction has been the subject of intense research and appears to be dependent on movement of the two to three intracellular loop of the DHPR which interacts with the RyR (140, 435). The RyR is a large and multiregulated protein potentially affected by many of the intracellular changes associated with fatigue (see sect. vii). It is known that SR  $\text{Ca}^{2+}$  release fails in various types of fatigue, and there is an unresolved debate as to whether this occurs through changes in the degree of voltage sensor activation or through the influence of changing myoplasmic metabolites or through depletion of  $\text{Ca}^{2+}$  inside the SR (see sects. vii, A and C, and ix).

$\text{Ca}^{2+}$  released by the SR gives rise to a transient increase in myoplasmic free  $[\text{Ca}^{2+}]$  ( $[\text{Ca}^{2+}]_i$ ), which is relatively easy to measure and can be regarded as the end product of the preceding processes.  $\text{Ca}^{2+}$  within the myoplasm binds to troponin C, instigates movement of tropomyosin, and allows the cycling of cross bridges, which eventually result in force development (19). The magnitude of the  $[\text{Ca}^{2+}]_i$  transients depends on the SR  $\text{Ca}^{2+}$  release and all the  $\text{Ca}^{2+}$  buffers in the cell, which include troponin C, parvalbumin, the SR  $\text{Ca}^{2+}$  pump, calmodulin,



and ATP (36). There are no measurements of  $\text{Ca}^{2+}$  buffering in fatigue which could assess how the  $\text{Ca}^{2+}$  buffering is affected by the ionic and metabolic changes of fatigue; consequently, the common assumption that changes in the  $[\text{Ca}^{2+}]_i$  transient reflect changes in SR  $\text{Ca}^{2+}$  release may be incorrect.

The sequence of EC coupling ends with cross-bridge activation. Cross-bridge properties can be determined in skinned fiber studies which allow the maximum  $\text{Ca}^{2+}$ -activated force ( $F_{\text{Ca,max}}$ ) and the  $\text{Ca}^{2+}$  sensitivity to be measured under conditions that simulate any chosen aspect of fatigue. Finally, the muscle relaxes as the elevated  $\text{Ca}^{2+}$  is pumped back into the SR by the ATP-driven SR  $\text{Ca}^{2+}$  pumps. The SR pumps are sensitive to many of the metabolic and ionic changes in fatigue, but the contribution of the changing pump properties to the slowing of relaxation in fatigue is still uncertain (8) (see sect. xi).

### III. METABOLIC CHANGES IN WORKING MUSCLES

A feature of fast muscle is that it can consume ATP, producing ADP and  $\text{P}_i$ , much faster than it regenerates it. Because the creatine kinase ( $\text{PCr} + \text{ADP} \leftrightarrow \text{Cr} + \text{ATP}$ ) and the adenylate kinase ( $2\text{ADP} \leftrightarrow \text{AMP} + \text{ATP}$ ) reactions are close to equilibrium, the net consumption of ATP leads to relatively stereotyped changes in the concentrations of ATP, ADP,  $\text{P}_i$ , phosphocreatine (PCr), creatine (Cr), and AMP which can be calculated from the equilibrium constants (10, 85). In essence, during net consumption of ATP, [ATP] is initially unchanged, and the net effect is a fall in [PCr] and rises in [Cr] and  $[\text{P}_i]$ , which have been observed in numerous biopsy and NMR studies of fatiguing muscles (e.g., Ref. 78). Later, when [PCr] reaches low levels ( $<10$  mM), [ATP] starts to fall and [ADP], which under control conditions would be around  $10 \mu\text{M}$ , rises substantially ( $100$ – $300 \mu\text{M}$ ). When [ADP] reaches such levels, the [AMP] also becomes significant and can be broken down by AMP deaminase to  $\text{NH}_3$  and inosine monophosphate (IMP).

Many studies have reported that cytoplasmic [ATP] does not drop below  $\sim 60\%$  of the resting level during either imposed stimulation or voluntary exercise (30, 117, 168, 286, 406, 455). However, virtually all of these measurements were made in whole muscle or muscle homogenates and hence reflect the spatially averaged change across all fibers present. Thus these values are difficult to interpret because the rate and extent of ATP usage differ substantially between fibers, depending on their metabolic profile and level of activation. If muscle biopsies are frozen and the individual fibers dissected out and typed, it becomes possible to determine the metabolic changes in identified fiber types. In a study of maximal cycling exercise in humans over 25 s, Karatzaferi et al. (238) showed

that when PCr was reduced to 11% ( $\sim 2.5$  mM) in the fastest fibers (type IIX), ATP was reduced to 20% ( $\sim 1.2$  mM) and IMP reached  $\sim 5$  mM. These changes are much greater than the mean changes across all the fiber types and occurred at a time when the power output had dropped to  $\sim 50\%$ . These experiments raise the possibility that changes in [ATP] may be more important than hitherto realized in intense exercise (see sect. viiC).

In addition to variations between fiber types, there must also be gradients of metabolites across the cells reflecting the different sites of consumption of ATP compared with its resynthesis. Calculations of the consumption and diffusion across myofibrils (assuming that ATP is consumed in the myofibrils and synthesized at mitochondria surrounding the myofibrils) indicate that the gradients of ATP are generally very small (311). Nevertheless, there may be sites in the cell where diffusion is very restricted, for instance, by multiple binding sites, where high local rates of ATP consumption lead to local depletion (197, 247, 453, 459).

One possible site of localized ATP depletion is the space between the t tubule and the SR (triad junction). ATP consumption in this region is substantial, owing to the presence of calcium pumps on the SR terminal cisternae just outside the junction (173, 392) and  $\text{Na}^+$ - $\text{K}^+$  pumps and other ATPases in the T-system membrane (206). Approximately 50% of all  $\text{Na}^+$ - $\text{K}^+$  pumps are in the T system (334), and the T system forms triad junctions with the SR for  $>90\%$  of its length in mammalian muscle fibers (138). Glycolytic enzymes associated with the triad junction support localized synthesis of ATP, which is used preferentially over cytoplasmic ATP (197). The glycolytic enzymes are well placed to utilize glucose entering the fiber via the T system (276), as well as the glucose-6-phosphate from adjacent glycogen stores.  $\text{Na}^+$ - $\text{K}^+$  pumps in muscle fibers preferentially use ATP derived from glycolysis (95, 337), including those pumps located in the T system (146). In view of the high density of ATP-consuming and -generating processes in the vicinity of the triad junction, as well as the comparatively small percentage of the cell volume it encompasses, [ATP] in the triad junction quite likely differs considerably from that in the cytoplasm as a whole. Thus this could allow the triad region to play a major role in sensing and responding to changes in cellular energy status, particularly given that the triad junction is the key transduction zone regulating  $\text{Ca}^{2+}$  release and contraction (see sect. viiC3).

When ADP is elevated, it is further hydrolyzed by the adenylate kinase reaction to AMP, which is then rapidly deaminated to IMP. This helps reduce the rise in [ADP] and consequent decrease in free energy for ATP hydrolysis. Most ADP in a resting muscle is bound to F-actin and does not directly influence metabolic reactions (453). During intense exercise, average cytoplasmic free [ADP] rises from  $\sim 10$  to  $\sim 200 \mu\text{M}$  (168, 455), and reductions in

[ATP] are nearly matched by equimolar rises in [IMP] (168, 238, 323, 407; see sect. x).

Another important factor is that when [ATP] decreases, free  $[Mg^{2+}]$  rises, because ADP, AMP, and IMP all have much lower affinity for  $Mg^{2+}$  than does ATP (55, 464). The free  $[Mg^{2+}]$  in resting muscle fibers is  $\sim 1$  mM (246, 464, 467). In single mouse fast-twitch fibers stimulated to fatigue, free  $[Mg^{2+}]$  rose rapidly at the time that force declined steeply and reached  $\sim 2$  mM when force had declined to  $\sim 30\%$  (464) (see sect. viiC3).

The pathways that resynthesize ATP include anaerobic glycogenolysis and the aerobic breakdown either of glycogen, glucose, or fat. Much current research focuses on the regulation of these pathways in different types of exercise (201, 402, 416) and is beyond the scope of this review. Anaerobic glycolysis is of central importance in muscle fatigue because it is turned on rapidly during activity, and the net reaction is breakdown of glucose units to lactate ions and protons causing the early acidosis associated with rapid-onset muscle fatigue. Typically this can lead to an acidosis of  $\sim 0.5$  pH units developing over several minutes at the start of intense exercise (for review, see Ref. 168). However, this acidosis is not an invariable accompaniment of muscle activity and presumably depends on the extent to which anaerobic glycolysis is switched on and the rate at which protons leave the muscle on the lactate transporter or the  $Na^+/H^+$  exchanger. Typically after long-lasting activity, particularly at a low duty cycle, there is little or no acidosis (91, 452) (see sect. viiB).

Glycogen is the principle store of energy in muscle and is rapidly depleted during intense exercise by anaerobic glycogenolysis and more slowly depleted in aerobic exercise. The twin correlations between muscle glycogen levels and a carbohydrate-rich diet and between glycogen depletion and exhaustion are two of the fundamental findings in exercise science (46); nevertheless, the reason why muscle fatigue is associated with glycogen depletion is still not understood (see sect. viiD).

#### IV. DIFFERENCES BETWEEN FAST AND SLOW FIBER TYPES

There are large differences between skeletal muscle cells regarding their speed of contraction, intracellular  $Ca^{2+}$  handling (see sect. ix), glycolytic versus oxidative capacity, fatigue resistance, etc. Several systems have been used to divide muscle cells into different fiber types, and the classification systems have to a large extent been a consequence of the methods available to distinguish between muscle cells (61, 410). At present, the dominating fiber type classification system for mammalian skeletal muscle is based on the expression of myosin heavy chain (MHC) isoforms, and the major fiber types are type

I, IIa, IIx, and IIb. While all four types of MHC are expressed in rodent muscles, IIb MHC is not expressed in human muscle (405). There are also muscle fibers that coexpress different MHC isoforms (354). The MHC isoform determines the rate of cross-bridge cycling and hence the maximal shortening velocity of a muscle cell, with type I being the slowest, type IIa intermediate, and IIx/b the fastest (61). However, it should be noted that the maximal shortening velocity differs between cells with the same MHC expression, which shows that it also depends on other factors, including the myosin light chain composition (60). In addition to MHC, there are isoforms of numerous other proteins with expression patterns that differ between muscle cells (61). In many instances, there is a distinguishable pattern of gene coexpression in cells so that the slow MHC type I is coexpressed with "slow" isoforms of other proteins. However, this is not always the situation, and there are multiple mechanisms interacting to control the gene expression of different protein isoforms (410).

A fiber type classification based on MHC isoforms has some relevance in the context of fatigue related to metabolic changes, because a fast isoform will consume ATP at a faster rate than a slow isoform. This can be illustrated by a recent study on isolated mouse fast-twitch fibers, where inhibition of cross-bridge cycling with *N*-benzyl-*p*-toluene sulfonamide (BTS) markedly delayed the fatigue-induced decrease in tetanic  $[Ca^{2+}]_i$  (66). Thus fatigue occurs more slowly when cross-bridge ATP consumption is decreased. However, a similar argument could be made for the other major ATP-consuming proteins in skeletal muscles, the SR  $Ca^{2+}$  pumps. These pumps exist in two isoforms, SERCA1 in fast type II fibers and SERCA2 in slow type I fibers (293), and the density of pumps is much higher in fast than in slow fibers (159). Furthermore, the major determinant of fatigue resistance is probably the muscle fiber's density of mitochondria and capacity to use oxidative metabolism. Slow type I fibers generally have a higher oxidative capacity than fast type II fibers (158), but in rat muscles, a higher oxidative capacity in type IIa than in type I fibers has been observed (24). In conclusion, "It is now clear that the MHC classification system for fiber types is a MHC, not a fiber type, classification system" (410). In relation to fatigue, slow type I fibers are generally more fatigue resistant than fast type II fibers, but this is mainly due to factors other than their MHC isoform.

#### V. HOW TO STUDY FATIGUE?

##### A. Models of Fatigue and Their Limitation

The "gold standard" for fatigue is the intact perfused muscle under central control. For selected muscle

groups, it is easy to measure the decline of force, and muscles can be studied by electromyography (EMG), NMR, and biopsy. A difficulty with each of these approaches is that muscles are mixtures of fiber types with very different properties, so there are problems either with the representativeness of the sample or the way in which spatial and/or temporal averaging disguises the interpretation. With EMG, it is difficult to distinguish the contributions from slowing of cortical firing rate, dispersion of conduction velocities along the fibers, change in resting potential, and AP amplitude (53). As noted in section 1, central fatigue is always a contributor to fatigue in intact animals. A useful variant of this approach is to produce fatigue by stimulating the nerve or muscle directly, thereby eliminating central contribution, and, because the blood flow is intact, eliminating the diffusion problems associated with isolated muscles.

Many studies of fatigue have been performed on isolated whole muscles, stimulated and perfused in a muscle bath. Because of the absence of the circulation, such preparations inevitably develop diffusion gradients of  $O_2$ ,  $K^+$ , and other substances across the muscle, and the apparent mechanisms of fatigue are likely to be biased towards these mechanisms. Barclay (32) has calculated the diffusion gradient of  $O_2$  across isolated mouse or rat muscles and concluded that an anoxic core is likely to develop whenever these preparations are repetitively activated. For instance, a whole mouse soleus can only contract at a duty cycle of 0.5 for  $\sim 60$  s at  $20^\circ C$  before an anoxic core develops. At  $35^\circ C$ , the anoxic core develops after only 12 s.  $K^+$  is released by active muscle fibers and accumulates in the extracellular space. It will therefore accumulate until a diffusion gradient develops sufficient to allow  $K^+$  to diffuse out of the preparation. Consequently, the concentration of extracellular  $K^+$  will be substantially higher in the center of the muscle than in the perfusate.  $CO_2$  and  $H^+$  and lactate will all tend to accumulate in the extracellular space in the same way. The combination of these issues makes analysis of mechanisms of fatigue in isolated whole muscles problematic. The relative contribution of extracellular accumulation of  $K^+$  or lactic acid or hypoxia was recently studied in isolated mouse muscle fatigued by repeated tetani, and under these experimental conditions, hypoxia was the most important factor (493).

Single fibers dissected from whole muscles eliminate the problems of  $O_2$  supply and the extracellular gradients of  $K^+$  and other products. In addition, there is only one fiber type under investigation, although identifying the fiber type is not trivial. An important additional advantage is the ease of performing fluorescent studies on single fibers. One disadvantage of this preparation is the considerable difficulty of learning the dissection. The absence of changes in  $O_2$  and  $K^+$  can also be viewed as a





disadvantage, since in the intact muscle changes in these substances clearly do occur and may contribute to fatigue. Typically single fibers are stimulated with plate electrodes so that the AP is simultaneously generated at multiple points along the fiber, eliminating longitudinal transmission of the AP as a possible mechanism of fatigue. Prolonged high-frequency stimulation applied focally to only one region of a muscle fiber produces greater and faster loss of force than does stimulation applied all along the length of the fiber via parallel electrodes (80, 265), which may indicate that failure of longitudinal AP transmission is an important fatigue mechanism in this situation. However, this force loss may reflect methodological problems with focal electrical stimulation in *in vitro* experiments, and in our view, this type of failure probably does not normally occur *in vivo* in healthy humans (see sect. vi) (see Table 1).

Single fibers do not survive well at  $37^\circ C$  or above (267), for reasons which are not entirely clear but seem to involve production of reactive oxygen species (315, 447). Thus a great deal of mammalian fatigue has been studied at unphysiological temperatures, an issue which is assuming increasing importance (see sect. vB).

Skinned fibers in which the surface membrane has been removed, either chemically or mechanically, have been widely used to study fatigue. Their main advantage is that the intracellular solution can be completely specified, and it is possible to examine the effects of changing individual metabolites rather than the range of changes that occur in intact fatiguing preparations. Traditionally skinned fibers were used to examine the response of the contractile proteins characterized by the  $[Ca^{2+}]$ -force relation. However, they can also be used to study SR  $Ca^{2+}$  release and uptake using the contractile proteins or an introduced  $[Ca^{2+}]$  indicator as a detector. Recently, it has been demonstrated that APs can be generated in the sealed off t tubules of mechanically skinned fibers (361), which means that the voltage sensors and the SR  $Ca^{2+}$  release channel and their interaction are amenable to experimental study. Disadvantages of skinned fibers are that it can be unclear whether a protein is present or absent in the preparation and the loss of soluble kinases, for instance, may have important effects on muscle function. In addition, metabolites whose importance has not been recognized by other means are generally not studied.

A great many subcellular components have also been studied as part of attempts to understand muscle function during fatigue. For instance, SR vesicles have been isolated from muscles in an attempt to understand the role of changing SR function in fatigue. Pump function of isolated SR vesicles can be determined, and in principle, such methods can be used to study how changing metabolites affect function or, by using standard conditions, how SR function has been changed by the fatigue process,

Table 1. *Advantages and disadvantages of various approaches to the study of fatigue*

|   |               |   |
|---|---------------|---|
|    | Advantages    | All physiological mechanisms present<br>Fatigue can be central or peripheral<br>All types of fatigue can be studied<br>Stimulation patterns appropriate for fiber types and stage of fatigue  |
|   | Disadvantages | Mixture of fiber types<br>Complex activation patterns<br>Produces correlative data; hard to identify mechanisms<br>Experimental interventions very limited  |
|    | Advantages    | Central fatigue eliminated<br>Dissection simple   |
|   | Disadvantages | Mixture of fiber types<br>Inevitable extracellular gradients of O <sub>2</sub> , CO <sub>2</sub> , K <sup>+</sup> , lactic acid<br>Mechanisms of fatigue biased by presence of extracellular gradients<br>Drugs cannot be applied rapidly because of diffusion gradients    |
|    | Advantages    | Only one fiber type present<br>Force and other changes (ionic, metabolic) can be unequivocally correlated<br>Fluorescent measurements of ions, metabolites, membrane potential, etc. possible<br>Easy and rapid application of extracellular drugs, ions, metabolites, etc. |
|   | Disadvantages | Dissection difficult<br>Environment different to in vivo<br>K <sup>+</sup> accumulation and other in vivo changes absent<br>Prone to damage at physiological temperatures<br>Small size makes analysis of metabolites difficult   |
|  | Advantages    | Precise solutions can be applied<br>Possible to study myofibrillar properties, SR release and uptake, AP/Ca <sup>2+</sup> release coupling<br>Metabolic and ionic changes associated with fatigue can be studied in isolation   |
|   | Disadvantages | Relevance to fatigue can be questionable<br>May lose important intracellular constituents<br>Relevant metabolites to study must be identified in other systems  |

assuming that the change is unaffected by isolation. SR release is difficult to study by such methods because the methods of stimulation of release (Ag<sup>+</sup>, caffeine) produce a rate of release slower by several orders of magnitude than the physiological mechanism. As in other areas of biology, studies of isolated elements of a cell can be highly effective at defining mechanisms but are generally incapable of demonstrating physiological relevance.

## B. Muscle Temperature

Several of the mechanisms that contribute to fatigue (e.g., the effects of intracellular pH and P<sub>i</sub> on contractile proteins, reactive oxygen species production) are temperature sensitive (see sects. VII, A and B, and VIII A). Furthermore, fatigue or decline in performance may occur more rapidly at high temperatures compared with low (121, 131, 315). For these reasons, it is important to know the temperature at which muscles normally operate. Muscle temperature is dependent on many factors including activity, blood flow, core temperature, closeness to body surface,

and environmental temperature. To eliminate this variability, it is common in studies of peripheral muscles in humans, e.g., adductor pollicis, to preheat the hand in a water bath at 45°C for 20 min, which sets the temperature of a superficial muscle to 37°C (121). The same approach can be used to set peripheral muscle temperature to any chosen temperature between 22 and 37°C (121). If whole body cooling is used to reduce both core and muscle temperature, muscle temperature may start as low as 30°C and increase to 38°C over a 35-min period of activity. This increase arises as a consequence of increased blood flow, increased core temperature, and heat generation by the muscle (58). Maximum muscle activity alone, in the absence of blood flow, causes a temperature rise of ~0.4°C/min (153). Studies in the mouse have shown that the temperature in the subcutaneous space of the foot was ~30°C (71); presumably during activity the muscles will warm up so that, as in humans, the range of temperature over which a muscle operates during a period of activity may be considerable.

In individuals performing intense activity in hot environmental conditions, it is widely appreciated that ex-



ercise performance declines for a variety of reasons that include muscle temperature, core temperature, and increased competition for blood flow to the muscle (131). Heat exhaustion in humans seems to occur at a relatively constant core temperature of  $\sim 40^{\circ}\text{C}$  (189), and despite a muscle temperature of  $40.8^{\circ}\text{C}$  (189), the failure of muscle performance in hyperthermia appears to be mainly central (439).

## VI. EXCITABILITY AND EXTRACELLULAR $\text{K}^+$ ACCUMULATION

### A. Basis of Excitability

#### 1. Spread of excitation

As mentioned in section II, normal muscle contraction depends on electrical excitation of the muscle fiber. Provided the muscle fiber is properly polarized, neuromuscular transmission initiates an AP, which propagates very rapidly along the muscle fiber surface (sarcolemma) in both directions (at  $\sim 2$  to  $6$  m/s in humans) (57) and then much more slowly ( $< 0.3$  m/s) (325) throughout the t-tubular system. The rapid sarcolemmal propagation is necessary to synchronously activate all parts of the muscle fiber to produce a useful contraction, and the conduction in the tubular system can be much slower because of the small distances involved (fiber diameter typically  $< 100\ \mu\text{m}$ ). It is necessary to have AP propagation in tubular system because passive spread of surface excitation into the T system is not enough to properly activate the voltage sensors and trigger  $\text{Ca}^{2+}$  release and contraction (5, 47, 148, 479). Most of the t-tubular system is oriented transversely, but some longitudinally oriented tubules are also present (172, 273). APs can also spread longitudinally inside the muscle fibers via these connections (253, 361), which helps ensure APs normally reach and stimulate every part of the t-tubular system even if conduction in some t tubules is hindered. This may explain why the formation of vacuoles (localized enlargements) in the t-tubular system during exercise does not seem to have any major deleterious effect on force production in most circumstances (262, 271).

#### 2. Types and distribution of ion channels

To understand the changes in excitability that occur with activity, it is necessary to consider the types and locations of the various ion channels involved. In adult muscle, the sodium current is carried by a rapidly activating and inactivating voltage-dependent channel ( $\text{Nav}1.4$ ) (233). The density of these  $\text{Na}^+$  channels is higher in fast-twitch than in slow-twitch muscle and is approximately threefold higher near the neuromuscular junction than at the ends of the fiber (384), presumably to

provide a safety factor for AP initiation there. Although the total number of  $\text{Na}^+$  channels on the sarcolemma is similar to that in the T system (224), the fivefold larger membrane area of the T system (138, 178) means that the  $\text{Na}^+$  channel density in the t tubules is only  $\sim 20\%$  of the surface density. In broad agreement, modeling of the AP in frog muscle indicated that the limiting  $\text{Na}^+$  conductance in the T system is  $\sim 5\%$  of that of the sarcolemma (5). The large  $\text{Na}^+$  current at the sarcolemma enables fast conduction and increases the likelihood of initiating an AP in each t tubule, where the relatively small tubular  $\text{Na}^+$  current is normally adequate for propagation (253).

Most of the resting  $\text{K}^+$  conductance is due to the  $\text{K}^+$  inward rectifier channels ( $\text{Kir}2.1$ ), which are present at higher density in the T system than at the sarcolemma in mammalian muscle (249). The most common potassium channel on the sarcolemma is the ATP-sensitive  $\text{K}^+$  channel ( $\text{K}_{\text{ATP}}$  or  $\text{Kir}6.2$ ) (417); these channels are also found in the T system but at a lower density (331). There are also  $\text{Ca}^{2+}$ -activated large conductance  $\text{K}^+$  channels (referred to as  $\text{B}_\text{K}$  or  $\text{K}_{\text{Ca}1.1}$ ) at approximately equal density in the sarcolemma and T system (331).

Importantly, normal adult skeletal muscle has a high density of the  $\text{ClC}1$  chloride channel (233). In mammalian muscle at rest, including in humans, the  $\text{Cl}^-$  permeability is approximately four to five times higher than the  $\text{K}^+$  permeability (62, 137, 171), and in frog muscle it is approximately two times higher (213). Most of this high  $\text{Cl}^-$  permeability arises from the T system (100, 137).

#### 3. Importance of membrane potential, $\text{Na}^+$ gradient, and leak conductances

Muscle fibers have to be well polarized to get adequate voltage-sensor activation of  $\text{Ca}^{2+}$  release. This is because chronic depolarization interferes with two key processes. First, the voltage-dependent  $\text{Na}^+$  channels become dysfunctional because of both increased “slow inactivation” (over the course of minutes) (385, 386) and “fast inactivation” (165), and they also require a greater depolarization to be opened (165). Second, the voltage sensors themselves become unresponsive (inactivated) (4, 378) so that even direct depolarization of the T system cannot activate full  $\text{Ca}^{2+}$  release (139, 378). A large proportion of the  $\text{Na}^+$  channels have to be inactivated for there to be complete AP failure, which occurs with prolonged depolarization to potentials less negative than approximately  $-60$  to  $-55$  mV in rat muscle (375) and approximately  $-50$  mV in frog muscle (265). The combined effect of voltage-sensor inactivation and AP failure reduces peak tetanic force in mammalian fibers by  $\sim 25\%$  at  $-60$  mV (82), due mostly to voltage-sensor inactivation (93, 352), and fully abolishes responses at approximately  $-55$  mV, due to AP failure.

AP failure can occur at more polarized potentials if the electrochemical gradient for  $\text{Na}^+$  is reduced, either by a decrease in external  $[\text{Na}^+]$ , as might occur in the T system during intense repetitive activation (47, 80, 227, 265), or by a rise in intracellular  $[\text{Na}^+]$  (335). AP failure can also occur if there is an increase in the permeability of the surface or T-system membranes to  $\text{K}^+$  or  $\text{Cl}^-$ , because an AP can only propagate if the inward  $\text{Na}^+$  current exceeds the leak currents carried by  $\text{K}^+$  and  $\text{Cl}^-$  enough to depolarize the adjacent membrane to threshold (423a). For example, repeated stimulation of a muscle fiber (166, 167, 193) can lower intracellular [ATP] sufficiently to activate ATP-sensitive  $\text{K}^+$  channels ( $\text{K}_{\text{ATP}}$ ) (116, 417) and greatly increase the  $\text{K}^+$  leak current, possibly reducing T-system excitability, even though the fiber is well polarized. As discussed later, there are a number of situations where  $\text{K}^+$  and  $\text{Cl}^-$  permeabilities may be modulated, either upwards or downwards, to decrease or enhance membrane excitability, respectively.

## B. Alterations in Excitability

### 1. Changes in electrochemical gradients with activity

There is a large body of literature showing that repeated activation of a muscle causes net  $\text{K}^+$  efflux (95, 212), increased extracellular  $[\text{K}^+]$  ( $[\text{K}^+]_{\text{o}}$ ) close to the muscle (232), and decreased intracellular  $[\text{K}^+]$  ( $[\text{K}^+]_{\text{i}}$ ) (403). This is due to the efflux of  $\text{K}^+$  occurring with each AP. In rested muscle cells, there is  $\sim 160$  mM  $\text{K}^+$  intracellularly and  $\sim 4$  mM extracellularly (399). Although only a very small amount of  $\text{K}^+$  leaves the muscle cell on each AP ( $\sim 2$  and  $10$   $\mu\text{M}$  in slow- and fast-twitch mammalian muscle, respectively) (95, 399), repeated activity can increase the extracellular  $[\text{K}^+]$  near the muscle fibers considerably, with it reaching close to  $\sim 9$  mM in well-perfused muscle working at high intensity and possibly exceeding  $10$  mM in localized regions (399). The problem of  $\text{K}^+$  accumulation is potentially much greater inside the t tubules (95, 168, 265, 399), because they have  $\sim 80\%$  of the total membrane surface area but their volume is only  $\sim 1\%$  of the total fiber volume (138, 178).

### 2. Failure of excitation in *in vitro* preparations and *in vivo* with imposed stimulation

In some circumstances, the changes in the electrochemical gradients for  $\text{K}^+$  occurring with repeated activation cause substantial membrane depolarization, failure of excitation, and a reduction in force responses. For example, repeated stimulation of whole muscles *in vitro* for 5–10 min results in a decrease of  $\sim 30\%$  or more in  $[\text{K}^+]_{\text{i}}$  and a depolarization of 10–18 mV in rat fast- and slow-twitch fibers and frog fibers (25, 230). The relatively

rapid recovery afterwards of  $[\text{K}^+]_{\text{i}}$ , membrane potential, and some force suggested that part ( $<30\%$ ) of the observed fatigue may have been attributable to inadequate excitation due to the  $[\text{K}^+]$  changes (25, 438). Similarly, continuous 60-Hz stimulation of rat fast-twitch muscles *in vitro* caused considerable  $\text{K}^+$  efflux from the muscles (and  $\text{Na}^+$  influx), with force declining  $\sim 80\%$  in 30 s in tandem with a similar decline in the compound surface AP (M-wave) (96). These findings indicate that in certain *in vitro* situations with imposed stimulation regimes, activity-induced changes in ion concentrations, particularly that of  $\text{K}^+$ , are not adequately compensated for by activity of the  $\text{Na}^+$ - $\text{K}^+$  pumps and other means and lead to reductions in muscle excitability and force.

Other experiments have shown that imposing continuous high-frequency stimulation on muscles *in vivo* in humans caused force to decline rapidly (227) and that this was associated with a reduction in the surface AP (52). Both force and the compound AP rapidly recovered substantially when the rate of stimulation was reduced (52, 227), suggesting that the force reduction was due to failure of excitation arising from the associated  $[\text{K}^+]$  changes and membrane depolarization.

In isolated single fibers *in vitro*, continuous high-frequency stimulation has also been shown to cause rapid force decline, fiber depolarization, and reductions in the intracellularly recorded AP (266). It was further observed that as tetanus duration increased, the cytoplasmic  $[\text{Ca}^{2+}]$  decreased in the middle of the fiber compared with the outer regions, indicating AP failure in the T system (148, 479). As force and cytoplasmic  $[\text{Ca}^{2+}]$  recovered when the stimulation frequency was reduced, the failure with high frequency of stimulation was likely due to fiber depolarization caused by excessive  $\text{K}^+$  accumulation in the T system, and possibly also to an additional effect on AP propagation caused by  $\text{Na}^+$  depletion in the T system. Voltage sensor inactivation probably did not contribute substantially to the reduced  $\text{Ca}^{2+}$  release, because experiments in muscle fiber bundles with comparable stimulation demonstrated that depolarization of the T system with high  $[\text{K}^+]$  solutions could still elicit a maximal response (81). Other findings in skinned fibers indicate that if a fiber is depolarized sufficiently to affect AP conduction in the T system, it interferes first with AP repriming, such that a single AP is able to propagate into and throughout the T system and elicit force, but subsequent closely spaced APs in a train cannot (147). Such AP failure is also indicated by findings in whole muscles *in vitro* where following repeated stimulation the response to tetanic stimulation was virtually no bigger than the twitch response to a single AP, and when the muscle was rested briefly, the response to tetanic stimulation recovered rapidly and proportionately much more than did the twitch response (438).

When the single fibers were subjected to intermittent rather than continuous stimulation, fatigue occurred much more slowly, and excitation-induced  $\text{Ca}^{2+}$  release was uniform through the fiber (148, 475, 479), indicating AP conduction in the T system. Some studies in frog single fibers, however, found evidence of T-system AP failure even with intermittent stimulation when tested at short muscle lengths (152, 180).

In summary, in certain circumstances AP-induced  $\text{K}^+$  movements do appear to reduce muscle excitability and force responses, such as when isolated muscles in vitro are given intense or prolonged stimulation or when all the motor units in a muscle in vivo are stimulated at a relatively high frequency. The occurrence of excitation failure depends greatly on the pattern and frequency of the imposed stimulation and is most marked with continuous high-frequency stimulation.

### 3. Effects on excitability of raising $[\text{K}^+]$ in vitro

It is also well-known that raising the  $[\text{K}^+]$  in the solution bathing a muscle in vitro can depolarize the fibers and depress excitability and force responses in a graded manner (82, 93, 213, 339, 341, 374), with effects being apparent above  $\sim 7$  mM  $\text{K}^+$  and with complete failure of tetanic force occurring at  $\sim 12$ – $13$  mM  $\text{K}^+$ . This depression of excitability and force can be reversed in many circumstances by stimulation of the  $\text{Na}^+$ - $\text{K}^+$  pump (95, 333, 340). Importantly, these depressive effects of raised  $[\text{K}^+]$  in the whole muscle experiments take tens of minutes to reach steady state (82). This is not simply due to the delay in  $\text{K}^+$  diffusion through the muscle but also reflects the very important role of  $\text{Cl}^-$  in muscle excitability. As detailed above, the sarcolemma and T-system membranes are more permeable to  $\text{Cl}^-$  than to  $\text{K}^+$ , and consequently, even though in the long term  $\text{Cl}^-$  will passively redistribute across the muscle membrane in accordance with the membrane potential determined largely by  $\text{K}^+$  (62), in the short term the depolarizing effect of any increase in extracellular  $[\text{K}^+]$  is greatly attenuated by the polarizing effects of  $\text{Cl}^-$  movement (83, 136, 213). Any consideration of the effects of activity-induced ion fluxes in muscle fibers in vivo needs to take in account 1) the magnitude and locality of the fluxes through the different ion channels and pumps; 2) the prevailing electrochemical gradients for  $\text{Na}^+$ ,  $\text{K}^+$ , and also  $\text{Cl}^-$ ; and 3) the intensity and duration of the exercise or stimulation. The lack of effect of  $[\text{K}^+]_o$  changes in the short term is likely to be indicative of what happens in vivo early in exercise, and the depressive effect seen in the longer term is possibly more relevant to the effects of very prolonged exercise when  $[\text{Cl}^-]_i$  may also change and the fibers become substantially depolarized.

### 4. Does reduced excitability cause muscle fatigue during normal exercise?

The key question then is whether or to what extent muscle excitability decreases in exercising muscles in vivo. A number of studies have examined what happens to the compound surface AP (M-wave) over the course of fatiguing exercising in humans. With continuous maximal voluntary contraction (MVC) of the adductor pollicis muscle for 60 s, force declined by  $\sim 40$ – $60\%$ , but the M-wave evoked by a single supramaximal stimulus was increased not decreased (52), indicating that excitation had not failed, at least not at the sarcolemma. It was also shown that direct stimulation of the muscle at various times points produced the same force as voluntary contraction (51). Sustained contraction of the quadriceps at 30% MVC for 3 min, sufficient to reduce twitch response by  $>50\%$  and raise venous plasma  $[\text{K}^+]$  from 4 to 6 mM, also caused no change in the M-wave (461). Similarly, with repeated intermittent contractions of the adductor pollicis muscle at 50% MVC until exhaustion ( $\sim 5$  min), there was no reduction in the M-wave, even though MVC and the evoked force response to 50 Hz direct stimulation were both decreased by  $\sim 50\%$  and the twitch response was decreased by  $\sim 70\%$  (22, 48, 49). A recent study with cycling exercise also found no change in the M-wave at the point of fatigue (394), though some other studies from the same group have observed some decrement. Thus reduced sarcolemma excitability was not responsible for the muscle fatigue occurring during the exercise in humans in the above cases.

Furthermore, findings of in vitro experiments (see sect. viB2) suggest that excitability problems in the T system would be expected to reduce tetanic force proportionately more than twitch force, but this is not what is observed in exercising humans. In view of this, it seems that the muscle fatigue experienced by the subjects in these exercise studies was not due to problems with excitation of either the sarcolemma or T system but instead caused by “metabolic” changes occurring inside the muscle fibers. As discussed in section vii, the relatively reduced twitch response is likely to be due to reduction in  $\text{Ca}^{2+}$  release from the SR and changes in  $\text{Ca}^{2+}$  sensitivity and maximum force production of the contractile apparatus ( $F_{\text{Ca,max}}$ ).

## C. Factors Helping to Prevent Loss of Excitation During Normal Exercise

The lack of excitability problems during normal exercise may seem surprising. However, there are a large number of factors acting in concert to help reduce or prevent loss of muscle excitation during normal exercise in vivo. These work in large part by reducing the extent of the  $[\text{K}^+]_o$  increase or its depolarizing effect. Some of



these mechanisms aid in limiting the number of APs that a fiber undergoes, whereas others involve utilizing the particular ion pathways present in a muscle fiber.

### 1. Motor unit recruitment

During submaximal force production *in vivo*, the central nervous system can vary the motor units activated, so as to spread the work load across the pool of motor units and hence reduce the requirements on a given motor unit (157). Less fatiguable motor units can be utilized at lower work loads, and the highest power and most fatiguable units recruited only during very demanding tasks (48). Recruitment of additional motor units explains the increase in EMG occurring when subjects sustain force at a given submaximal level. In fact, the discharge rate of originally active motor units can decrease during prolonged submaximal contractions, despite the fact the overall excitatory drive to the motor neuron pool increases as fatigue develops (181).

### 2. Activation at optimal firing rate

Fast-twitch and slow-twitch motor units are normally activated *in vivo* at the firing rate most appropriate to the contractile properties of the motor unit (e.g., at ~30 and 10 Hz, respectively, in humans), such that it is just sufficient to give a fused tetanus of close to maximum force (42, 53). This keeps the number of APs used to excite a muscle fiber to the minimum necessary. Stimulating motor units at higher than their optimal firing rate, as may happen with imposed nerve or muscle stimulation, causes the muscle fibers to fatigue more rapidly than during voluntary contraction (227).

### 3. Decreasing firing rate

Motor neurons decrease their firing rate during sustained maximal contractions (50). This reduction is normally well-matched to the slowing of relaxation that occurs with muscle activation (see sect. xi), such that the stimulation rate remains just sufficient for eliciting the greatest tetanic force possible at that point in time (28, 51, 53). This phenomenon is often called "muscle wisdom" (157, 179). In addition, impulse trains *in vivo* typically begin with groups of closely spaced impulses (doublets), which appear to enable force production with less associated fatigue (54), likely because of the force potentiation occurring with the closely spaced impulses (1), which gives more effective force production for a given number of APs.

### 4. Sarcolemma AP changes

Repeated activation can slow the propagation of the surface AP (231), reduce its size, and increase its duration (28, 265), presumably owing to fiber depolarization. Nev-

ertheless, these changes to the surface AP do not appear to contribute to fatigue (28), as the AP evidently remains sufficient to initiate an AP in the T system because of the large safety factor in the process. The changes in the sarcolemmal properties and AP can also lead to intermittent failure in a train of APs (28, 80), but the accompanying slowing of fusion frequency means that such AP failure does not evidently contribute to decreased force.

### 5. $\text{Na}^+$ - $\text{K}^+$ pumping

The  $\text{Na}^+$ - $\text{K}^+$  pump plays a vital role in lowering  $[\text{K}^+]_o$  (95), particularly in the T system.  $\text{Na}^+$ - $\text{K}^+$  pump activity is stimulated by muscle excitation, raised intracellular  $[\text{Na}^+]$ ,  $\beta$ -adrenergic stimulation, and raised temperature (335, 341, 350) and can be increased more than 20-fold above its resting rate. In rat fast-twitch muscle, ~50% of all the  $\alpha_2$ -isoform of the pump are in the T system and are estimated to be able to clear  $\text{K}^+$  from there at ~4 mM/s (334). In addition, activity of the pump is electrogenic and can hyperpolarize a muscle fiber a further ~8 mV (95).

### 6. Important role of $\text{Cl}^-$ channels

The high resting permeability of the T system to chloride relative to potassium ( $P_{\text{Cl}}/P_{\text{K}} \sim 4$ ) (100, 137) means that the resting potential ( $E_{\text{M}}$ ) will be heavily weighted towards the chloride equilibrium potential ( $E_{\text{Cl}}$ ) (83, 213), which will remain relatively constant during activity owing to the high  $[\text{Cl}^-]$  in the T system (~110 mM) and relatively large amount of  $\text{Cl}^-$  influx needed to change the  $[\text{Cl}^-]_i$ . For this reason, any inward flux of  $\text{Cl}^-$  during the repolarizing phase of an AP has much less effect on steady-state conditions than would a matching outward flux of  $\text{K}^+$  (62, 137).

Importantly too, the high  $\text{Cl}^-$  permeability of the T system likely aids in recovering some of the  $\text{K}^+$  that does build-up in the T system, because if the  $[\text{K}^+]$  in the T system increases sufficiently such that  $E_{\text{K}}$  across the T-system membrane is less negative than  $E_{\text{M}}$ ,  $\text{K}^+$  will be driven by their electrochemical gradient back into the fiber through the inward rectifier channels. The importance of this effect has been indicated by quantitative modeling (454), though it should be noted that the Wallinga et al. (454) study considerably underestimated the magnitude of the effect as it assumed the T-system chloride conductance was almost 10-fold lower than its true value (62, 100, 137). Thus the high chloride conductance in the T system may not only reduce the rate of accumulation of  $\text{K}^+$  there, but also help the recovery of  $\text{K}^+$ , greatly reducing the problem of  $\text{K}^+$  movements on membrane potential and excitability.

The importance of the  $\text{Cl}^-$  conductance in mammalian muscle is apparent from the problems that occur with muscle function when it is absent (3, 62, 445). Individuals



with generalized (or Bekker's) myotonia, where there is no muscle  $\text{Cl}^-$  conductance, are unable to maintain a 60% MVC, and force drops within a few seconds, associated with progressive then complete failure of the sarcolemma AP in most fibers. This effect likely occurs because the high frequency of APs causes a rapid build-up of  $\text{K}^+$  in the T system, which depolarizes the fiber as a whole, and causes sarcolemma AP failure (445). Interestingly, the subjects are able to subsequently increase force output over the following 20–60 s to close to 60% MVC again ("warm-up" phenomenon), which appears to be due to the reduced rate of AP stimulation of the muscle occurring at those later times (445).

### 7. Alterations in leak conductances

If fibers do become depolarized with repeated activity, progressively more of the  $\text{Na}^+$  channels will become inactivated. This effect can be beneficial, because the smaller  $\text{Na}^+$  influx will mean less  $\text{K}^+$  and  $\text{Cl}^-$  flux is needed for repolarization. However, AP propagation will fail if the  $\text{Na}^+$  current is not large enough to overcome the resting leak current, which is carried primarily by  $\text{Cl}^-$  in mammalian muscle. To help avoid this problem, muscle fibers have mechanisms for reducing the  $\text{Cl}^-$  leak conductance during activity. Increasing intracellular acidity (e.g., to pH 6.6), as can occur during heavy exercise, reduces the resting  $\text{Cl}^-$  conductance approximately two-fold, enabling APs to continue to generate large tetanic force even when a fiber is depolarized to close to  $-60$  mV (333, 351, 352).

### 8. $\text{Ca}^{2+}$ release feedback

A further factor helping maintain  $\text{Ca}^{2+}$  release is that the amount of  $\text{Ca}^{2+}$  released per AP is to some extent kept constant by feedback mechanisms operating on the AP- $\text{Ca}^{2+}$  release coupling. The amount of  $\text{Ca}^{2+}$  release triggered by an AP remains almost constant even if the amount of  $\text{Ca}^{2+}$  in the SR is increased greatly above its normal endogenous level (360) (see sect. 1xB and Fig. 5). This occurs because the  $\text{Ca}^{2+}$  released into the narrow junctional gap 1) causes  $\text{Ca}^{2+}$ -dependent inactivation of the SR release channel (35, 346, 398), 2) speeds the deactivation of the adjacent voltage sensors (345), and 3) shortens the duration of the AP by activating the  $\text{Ca}^{2+}$ -activated  $\text{K}^+$  channels and possibly  $\text{Ca}^{2+}$ -activated  $\text{Cl}^-$  channels in the T system (345). These three factors normally limit  $\text{Ca}^{2+}$  release (360). Consequently, if the T-system AP becomes smaller and/or the voltage sensors become less activated, resulting in a drop in the rate of SR  $\text{Ca}^{2+}$  release, the above inhibitory effects will be slower in onset, thus helping keep up the total amount of  $\text{Ca}^{2+}$  release per AP.

In summary, although many in vitro and in vivo studies show that a loss of muscle excitability can occur with

imposed stimulation, this does not usually appear to be the cause of fatigue in exercising humans. This is probably because many compensatory processes act together to prevent changes in  $[\text{K}^+]$  and fiber depolarization or to reduce their deleterious effects.

## VII. METABOLIC CHANGES AND FATIGUE

During locomotion and most other types of muscle activity, muscles are activated with repeated short bursts of APs. In the following section of this review we will therefore focus on models where fatigue has been induced by repeated short contractions. In fast-twitch fibers, this type of fatiguing stimulation results in a characteristic pattern of changes in force and  $[\text{Ca}^{2+}]_i$ . Initially, there is a fast decline of tetanic force by 10–20% that is accompanied by an increase in tetanic  $[\text{Ca}^{2+}]_i$  (phase 1), then follows a period of relatively constant tetanic force (phase 2); finally, there is rapid decline of both tetanic force and  $[\text{Ca}^{2+}]_i$  (phase 3) (Figs. 1 and 3A). The importance of the decline of tetanic  $[\text{Ca}^{2+}]_i$  in the phase 3 decline of force has been established by rapid application of caffeine at concentrations that increase the opening of the SR  $\text{Ca}^{2+}$  channel (149, 268, 462). Caffeine can cause recovery of the tetanic  $[\text{Ca}^{2+}]_i$  in phase 3 and reverse much of the final phase of force decline. However, this effect has not yet convincingly been demonstrated in intact, perfused muscles (218).

Interestingly, the pattern of force decline is similar in frog, mouse, and rat fast-twitch single fibers (9, 283, 287, 420, 462). A similar pattern of force decline was also described when single fast-twitch motor units were stimulated in anesthetized cats (75). The pattern with an initial increase in tetanic  $[\text{Ca}^{2+}]_i$  followed by a decline as fatiguing stimulation continues has also been observed in isolated whole mouse extensor digitorum longus (EDL) muscles (203), whereas tetanic  $[\text{Ca}^{2+}]_i$  decreased monotonically in bullfrog semitendinosus muscles (23).

The duration of phase 2 (i.e., the period with stable force production) differs markedly between cells, which seems to reflect differences in the ability to use oxidative metabolism, because it is markedly shortened by mitochondrial inhibition with cyanide (Fig. 1) (268, 462) or exposure to decreased oxygen pressure (420). Whole muscles generally do not show a clear phase 2, which probably reflects differences in fatigue resistance among the individual fibers within the muscle.

In the experiments described above, the stimulation frequency during the tetani was set close to that required to produce maximal tetanic force. This indicates that the initial force decrease during phase 1, where tetanic  $[\text{Ca}^{2+}]_i$  actually increases, is caused by a reduction in cross-bridge force-generating capacity. Conversely, the final rapid force decrease during phase 3 would be due to




FIG. 3. Tetanic  $[Ca^{2+}]_i$  and force records at various phases of fatigue obtained in a fast-twitch (A) and a slow-twitch (B) mouse fiber. Numbers above  $[Ca^{2+}]_i$  records indicate order of tetani. The fast-twitch FDB fiber in A shows the normal pattern of easily fatigued fibers: an early increase in tetanic  $[Ca^{2+}]_i$  accompanied by  $\sim 10\%$  decrease in force (phase 1; tetanus 1–10), followed by a relatively stable period (phase 2; tetanus 10–60), and finally a rapid decrease of tetanic  $[Ca^{2+}]_i$  and force (phase 3; tetanus 60–88). Stimulation protocol: 350-ms, 70-Hz tetani given at 2.5-s intervals. The fatigue-resistant soleus fiber in B shows little changes in tetanic  $[Ca^{2+}]_i$  and force during a markedly more demanding fatiguing stimulation protocol [a total of 1,000 tetani (500 ms, 70 Hz) given at 2-s intervals]. [A from Dahlstedt et al. (107); B from Bruton et al. (67).]

the combined effect of decreased tetanic  $[Ca^{2+}]_i$  (see Fig. 3A) and reduced myofibrillar  $Ca^{2+}$  sensitivity.

Isolated slow-twitch fibers and motor units are generally highly resistant to fatigue induced by repeated short tetani (67, 75, 188, 250). Changes in tetanic  $[Ca^{2+}]_i$  during fatigue of single slow-twitch fibers have been little studied. In mouse soleus fibers, tetanic  $[Ca^{2+}]_i$  followed the pattern observed in fast twitch fibers, i.e., an initial increase followed by a decrease (67) (Fig. 3B), whereas the early increase in tetanic  $[Ca^{2+}]_i$  was not observed in rat soleus fibers (288).

In the following sections we discuss how force is affected by metabolic changes that occur during fatigue. We focus on fast-twitch muscle fibers where metabolic changes generally are larger and the force decrease more marked.

### A. Inorganic Phosphate

The exchange of phosphate between ATP and PCr is catalyzed by creatine kinase (CK) according to the following reaction:  $PCr + ADP + H^+ \rightarrow Cr \text{ (creatine)} + ATP$ . During periods of high energy demand, the ATP concentration initially remains almost constant while CrP breaks down to Cr and  $P_i$ . While Cr has little effect on contractile function (321),  $P_i$  may cause a marked decrease of myofibrillar force production and  $Ca^{2+}$  sensitivity as well as SR  $Ca^{2+}$  release. Accordingly, increased  $P_i$  is considered to be a major cause of fatigue (470). A causative role of increased  $P_i$  has also been implied in other situations with impaired muscle function. For instance, a recent study where human subjects were followed during the rehabilitation after cast immobilization shows a significant in-

verse relationship between resting  $P_i$  and specific force production (349).

### 1. $P_i$ and cross-bridge force production

According to present models of cross-bridge force production, the myosin head is first bound weakly and then strongly to the actin filament. Thereafter  $P_i$  is released, possibly resulting in a further increase in force production (433). This implies that the transition to high-force cross-bridge states is inhibited by increased  $P_i$ , and fewer cross bridges would be in high-force states when  $P_i$  increases during fatiguing stimulation. In line with this, experiments on skinned fibers consistently show a substantial decrease of  $F_{Ca,max}$  in the presence of elevated  $P_i$  (314, 348). Mammalian skinned fiber experiments are generally performed at temperatures much lower than those prevailing *in vivo*, and the  $P_i$ -induced inhibition of cross-bridge force production becomes less marked as temperature is increased (101, 112, 123).

The effect of  $P_i$  on cross-bridge force production has been difficult to directly test in intact muscle cells, because it has proven difficult to alter myoplasmic  $P_i$  without imposing other metabolic changes as well. An experimental model that can be used in this regard is genetically modified mice that completely lack CK in their skeletal muscles (CK<sup>-/-</sup> mice) (422). Skeletal muscle fibers of CK<sup>-/-</sup> mice display an increased myoplasmic  $P_i$  concentration at rest, and there is no significant  $P_i$  accumulation during fatigue (422). The  $F_{Ca,max}$  of unfatigued CK<sup>-/-</sup> fast-twitch fibers is markedly lower than that of wild-type fibers, and this can partly be explained by a  $P_i$ -induced depression of cross-bridge force production (106). Furthermore, CK<sup>-/-</sup> fibers do not display the 10–20% reduction of  $F_{Ca,max}$  observed after ~10 fatiguing tetani in fast-twitch fibers (phase 1; see Figs. 1 and 3A), which has been ascribed to increased  $P_i$  (106, 107). Even after 100 fatiguing tetani, force was not significantly affected in CK<sup>-/-</sup> fibers, whereas by this time force was reduced to <30% of the original in wild-type fibers. Additional support for a coupling between myoplasmic  $P_i$  concentration and force production in intact muscle cells comes from experiments where reduced myoplasmic  $P_i$  is associated with increased force production (73, 74, 355).

In conclusion, increased myoplasmic  $P_i$  can inhibit force production by direct action on cross-bridge function, and this is a likely mechanism underlying the decrease in tetanic force occurring early during fatigue in fast-twitch fibers. The magnitude of this  $P_i$ -induced decrease in cross-bridge force production is probably rather small (~10% of maximum force) in mammalian muscle at physiological temperatures.

### 2. $P_i$ and myofibrillar $Ca^{2+}$ sensitivity

Altered cross-bridge function may also affect the force- $[Ca^{2+}]_i$  relationship via the complex interaction between myosin cross-bridge attachment and thin (actin) filament activation (192), and reduced myofibrillar  $Ca^{2+}$  sensitivity is frequently observed in skeletal muscle fatigue (168, 480). Skinned fiber experiments have shown that increased  $P_i$  decreases the myofibrillar  $Ca^{2+}$  sensitivity (301, 314). Intriguingly, the inhibitory effect of  $P_i$  on myofibrillar  $Ca^{2+}$  sensitivity was recently shown to be larger at 30 than at 15°C (124), i.e., opposite to the inhibitory effect of  $P_i$  on cross-bridge force production at saturating  $[Ca^{2+}]_i$ , which is smaller at high temperature (123). Results from unfatigued CK<sup>-/-</sup> fibers, which display an increased myoplasmic  $P_i$  concentration at rest, also indicate a  $P_i$ -induced decrease in the myofibrillar  $Ca^{2+}$  sensitivity (106).

In summary, a fatigue-induced increase in  $P_i$  can reduce myofibrillar  $Ca^{2+}$  sensitivity, which may have a large impact on force production in later stages of fatigue where tetanic  $[Ca^{2+}]_i$  decreases.

### 3. $P_i$ and the increase of tetanic $[Ca^{2+}]_i$ in early fatigue

An increase in tetanic  $[Ca^{2+}]_i$  is generally observed early during fatigue induced by repeated tetanic stimulation (phase 1, see above). This increase was not observed in CK<sup>-/-</sup> muscle fibers (106, 107), but reappeared after CK injection into these fibers (105). A similar result has been observed in *Xenopus* fibers when CK was blocked pharmacologically (243). These results indicate a key role of CK and increased  $P_i$  in the early increase in tetanic  $[Ca^{2+}]_i$ . There are several mechanisms by which changes in CK activity and increased  $P_i$  can lead to increased tetanic  $[Ca^{2+}]_i$ . First, increased  $P_i$  might reduce the myoplasmic  $Ca^{2+}$  buffering by decreased binding of  $Ca^{2+}$  to troponin C due to a reduction in strong cross-bridge attachment (314), but this mechanism appears to be of limited importance in skeletal muscle (192). Second,  $P_i$  might act on the RyR and increase SR  $Ca^{2+}$  release. Accordingly, elevated  $P_i$  has been shown to increase the open probability of isolated RyR incorporated into lipid bilayers as well as the rate of  $Ca^{2+}$ -induced  $Ca^{2+}$  release in SR vesicles and skinned fibers (27, 174). However, it has since been shown that  $P_i$  decreases caffeine- and depolarization-induced SR  $Ca^{2+}$  release in mechanically skinned rat skeletal muscle fibers in a  $Mg^{2+}$ -dependent manner (135). The reason for these conflicting results is not clear but is likely to involve differences in experimental conditions, e.g., mode of SR activation and the presence or absence of proteins associated with the RyR (135). Third, elevated  $P_i$  may inhibit (118, 426) or even reverse (133) SR  $Ca^{2+}$  pumping which, at least in the short term, might increase tetanic  $[Ca^{2+}]_i$  (468). In addition,

tion, removal of PCr inhibits SR  $\text{Ca}^{2+}$  uptake in skinned fibers (132), supposedly caused by decreased ATP buffering via the CK reaction leading to a decreased ATP/ADP· $\text{P}_i$  ratio in the vicinity of the SR  $\text{Ca}^{2+}$  pumps (134, 423). An important role of CK and  $\text{P}_i$  on SR  $\text{Ca}^{2+}$  pumping is supported by the finding that the early increase in tetanic  $[\text{Ca}^{2+}]_i$  during fatigue is accompanied by a marked increase in the amplitude of  $[\text{Ca}^{2+}]_i$  tails during relaxation in normal fibers but not in CK<sup>-/-</sup> fibers (106).

In conclusion, an operating CK system and  $\text{P}_i$  accumulation are required for the early increase in tetanic  $[\text{Ca}^{2+}]_i$  during fatigue, but the exact mechanism(s) involved remains to be established.

#### 4. $\text{P}_i$ and the decrease of tetanic $[\text{Ca}^{2+}]_i$ in late fatigue

After the initial increase, tetanic  $[\text{Ca}^{2+}]_i$  decreases as fatiguing stimulation progresses, and this, together with a decreased myofibrillar  $\text{Ca}^{2+}$  sensitivity, results in the final rapid decrease of tetanic force (phase 3). CK and elevated  $\text{P}_i$  appear to play a central role in this decrease because it is markedly delayed in fibers with CK inhibition induced either pharmacologically (108) or genetically (CK<sup>-/-</sup> muscles) (107). Moreover, CK injection into CK<sup>-/-</sup> fibers resulted in the normal changes in tetanic  $[\text{Ca}^{2+}]_i$  during fatigue, i.e., an early increase followed by a decrease (105). We will discuss two mechanisms by which elevated  $\text{P}_i$  may decrease SR  $\text{Ca}^{2+}$  release:  $\text{P}_i$ -induced inhibition of the RyR and  $\text{Ca}^{2+}$ - $\text{P}_i$  precipitation in the SR, causing a reduction of the amount of free  $\text{Ca}^{2+}$  available for release.

Skinned fiber experiments have shown an inhibitory effect of  $\text{P}_i$  on caffeine- and depolarization-induced SR  $\text{Ca}^{2+}$  release (135). This inhibition appears distinct from the  $\text{P}_i$  effect on SR  $\text{Ca}^{2+}$  pumping and  $\text{Ca}^{2+}$ - $\text{P}_i$  precipitation in the SR and is attributed to a  $\text{P}_i$  effect on the SR  $\text{Ca}^{2+}$  release mechanism (135, 423). Interestingly, this  $\text{P}_i$ -induced inhibition depends on changes in  $\text{Mg}^{2+}$  within the physiological range (135): the inhibitory effect was markedly larger at the free  $[\text{Mg}^{2+}]$  of the myoplasm of fatigued (~1.6 mM) than rested (~0.8 mM) mammalian muscle fibers (464). Thus this  $\text{Mg}^{2+}$ -dependent  $\text{P}_i$ -induced inhibition of SR  $\text{Ca}^{2+}$  release is consistent with an inverse relationship between tetanic  $[\text{Ca}^{2+}]_i$  and  $[\text{Mg}^{2+}]_i$  during fatigue under normal conditions, which is completely lost after pharmacological inhibition of CK (108).

$\text{Ca}^{2+}$ - $\text{P}_i$  precipitation in the SR may be an important mechanism behind the decline in tetanic  $[\text{Ca}^{2+}]_i$  in late stages of fatigue (11, 142, 423). The underlying theory is as follows: the free  $[\text{Ca}^{2+}]$  in the SR ( $[\text{Ca}^{2+}]_{\text{SR}}$ ) is assumed to be ~1 mM, and the  $\text{Ca}^{2+}$ - $\text{P}_i$  solubility product, measured in vitro, is ~6 mM<sup>2</sup> (175). When  $\text{P}_i$  increases in the myoplasm during fatigue, some  $\text{P}_i$  ions subsequently enter the SR, and when the  $\text{Ca}^{2+}$ - $\text{P}_i$  solubility product is exceeded, precipitation occurs and the  $[\text{Ca}^{2+}]_{\text{SR}}$  decreases (175,

222). This reduction in the releasable pool of  $\text{Ca}^{2+}$  in the SR can, depending on the initial SR  $\text{Ca}^{2+}$  load (360), lead to decreased SR  $\text{Ca}^{2+}$  release.

The occurrence of SR  $\text{Ca}^{2+}$ - $\text{P}_i$  precipitation has not been directly shown, but indirect evidence of its existence has been presented both in experiments on skinned fibers with intact SR exposed to high  $\text{P}_i$  solutions (135, 142, 175) and in intact mouse fibers injected with  $\text{P}_i$  (469). In the latter study,  $\text{P}_i$  was injected into unfatigued mouse fast-twitch muscle fibers, and this resulted in decreased tetanic  $[\text{Ca}^{2+}]_i$ , faster SR  $\text{Ca}^{2+}$  reuptake, and decreased resting  $[\text{Ca}^{2+}]_i$ , all of which are consistent with a decreased  $[\text{Ca}^{2+}]_{\text{SR}}$ . Several fatigue experiments also support the  $\text{Ca}^{2+}$ - $\text{P}_i$  precipitation mechanism. First,  $[\text{Ca}^{2+}]_i$  in response to a high dose of caffeine or 4-choro-*m*-cresol is decreased in fatigued mouse (462) and toad muscle fibers (234). These compounds directly stimulate the SR  $\text{Ca}^{2+}$  release channels, and a decreased response indicates a reduced pool of  $\text{Ca}^{2+}$  available for release. Second, the  $[\text{Ca}^{2+}]_{\text{SR}}$ , measured with the low-affinity  $\text{Ca}^{2+}$  indicator fluo-5N, declined throughout a period of fatiguing stimulation of toad muscle fibers and recovered afterwards (235). The recovery occurred even in the absence of extracellular  $\text{Ca}^{2+}$ , which shows that  $\text{Ca}^{2+}$  did not disappear from the cells during fatigue. On the other hand, recovery was blocked by mitochondrial inhibition with cyanide, showing that it depended on aerobic metabolism.

While there are numerous results supporting an important role of  $\text{Ca}^{2+}$ - $\text{P}_i$  precipitation in fatigue, there are also several results that are difficult to reconcile with this mechanism (423). For instance, the marked increase in  $\text{P}_i$  develops early during fatiguing stimulation, whereas the decrease in tetanic  $[\text{Ca}^{2+}]_i$  occurs with a delay. Moreover, in mouse fast-twitch fibers, the decline of tetanic  $[\text{Ca}^{2+}]_i$  temporally correlates with an increase in cytosolic  $[\text{Mg}^{2+}]$ , which presumably stems from a net breakdown of ATP (464), and it is not obvious why  $\text{Ca}^{2+}$ - $\text{P}_i$  precipitation in the SR should show a temporal correlation with increasing  $[\text{Mg}^{2+}]$ /decreasing  $[\text{ATP}]$ . However, the SR membrane contains small-conductance chloride channels, which may conduct  $\text{P}_i$  (6, 279). The open probability of these channels increases at low ATP, which fits the finding that  $\text{P}_i$  entry into the SR of skinned muscle fibers is inhibited by ATP (359), although others did not observe this ATP dependency (142, 423). Other uncertainties regarding the importance of SR  $\text{Ca}^{2+}$ - $\text{P}_i$  precipitation in fatigue include (423) the concentrations of  $\text{Ca}^{2+}$  and  $\text{P}_i$  required to cause precipitation might be higher in the functional SR than in simple salt solutions,  $\text{Ca}^{2+}$ - $\text{P}_i$  may exist in many different forms (456), and unstable complexes dissolve rapidly, which means that more  $\text{Ca}^{2+}$  might become available as the  $[\text{Ca}^{2+}]_{\text{SR}}$  declines during a tetanic contraction.



In conclusion, while it is clear that increased  $P_i$  can cause a decrease in tetanic  $[Ca^{2+}]_i$  in late fatigue, the importance of this in different types of fatigue and the exact processes involved require further attention.

### 5. Creatine supplementation

There is widespread use of extra Cr intake among athletes, not only in the elite but also among people exercising on a recreational level (for recent reviews, see Refs. 44, 437). Cr supplementation is also used in an attempt to improve muscle function in diseases such as inflammatory and mitochondrial myopathies and muscle dystrophies. Cr enters muscle cells via a  $Na^+$ -dependent transporter in the sarcolemma. Inside the cell, Cr is phosphorylated by CK, and the  $[PCr]/[Cr]$  ratio basically depends on the energy state of the cell. Extra Cr intake increases the total muscle Cr content by up to ~20%. Cr supplementation has a modest positive effect on muscle performance during bouts of short-term (~10 s) high-power exercise, whereas it does not improve function during more long-lasting types of muscle activity (437). This fits with the fact that PCr breakdown contributes to a relatively large fraction of the ATP supply during the first seconds of high-intensity muscle activity, whereas the PCr contribution is minimal during prolonged exercise.

On the muscle cell level, an increased PCr concentration will provide better ATP buffering during intense activity and delay increases in ADP that might slow cross-bridge cycling (99, 473) and SR  $Ca^{2+}$  pumping (132, 222). Thus a high power output could be sustained for a slightly longer time. Moreover, PCr/Cr is osmotically active, and short-term Cr supplementation is generally accompanied by increased body weight due to water accumulation in muscle cells (321, 437). The increased water content in muscle cells might improve function by increasing the myofibrillar  $Ca^{2+}$  sensitivity and the  $F_{Ca,max}$  (321).

In the context of fatigue at the cellular level, it might be expected that increased PCr loading in muscle cells causes a more marked decrease in contractile function during intense activity due to a larger increase in  $P_i$ , which is considered to be a major cause of fatigue inhibiting both myofibrillar and SR function (see above). However, it should be noted that cross-bridge force production and myofibrillar  $Ca^{2+}$  sensitivity decline linearly with the logarithm of  $[P_i]$  (314, 348); in other words, the depressive effect of an increase in  $P_i$  is large at the onset of fatigue where  $[P_i]$  is low, but it becomes more and more limited as fatigue progresses and  $[P_i]$  reaches high levels. The  $[P_i]$  dependency on the fatigue-induced changes in SR  $Ca^{2+}$  handling is probably complex, and the present understanding of the processes involved is too limited to predict the effect on these processes of increased PCr loading. It should be noted that the beneficial effect of

increased PCr is observed in short-lasting bouts of exercise where inhibitory effects of energy metabolites on SR  $Ca^{2+}$  release are limited (see above).

## B. Lactate and $H^+$

The accumulation of lactic acid in muscle has historically been suggested to be the major cause of muscle fatigue (see review in Ref. 168). As outlined in section III, lactate and  $H^+$  are produced in muscles during intense exercise (379), and in humans, the intracellular lactate concentration may reach 30 mM or more (expressed relative to cell water) and the intracellular pH ( $pH_i$ ) decreases by ~0.5 pH units (389). A close temporal relationship is often observed between decreased muscle force and increased intracellular concentrations of lactate and particularly  $H^+$ . However, such correlations break down in many cases, and although increased intracellular levels of  $H^+$  may reduce muscle performance to some degree, it now appears that its deleterious effects have been considerably overestimated and that other, beneficial, effects have been overlooked.

### 1. Lactate

In humans exercising at different work intensities, lactate levels were found not to correlate well with muscle fatigue (240). Addition of lactate to the outside of muscles in vivo (214) or in vitro (411) resulted in some reduction in tetanic force, but a large part of this effect was likely due not to lactate uptake but instead to the increased extracellular osmolality causing water to move out of the muscle fibers thereby increasing intracellular ionic strength, which has direct inhibitory effects on force production (259). Experiments with skinned muscle fibers at constant ionic strength have shown that lactate, at concentration even up to 50 mM, has relatively little effect on force production by the contractile apparatus, reducing  $F_{Ca,max}$  by <5% and having little or no significant effect on  $Ca^{2+}$  sensitivity (16, 86, 143, 358). Furthermore, even though lactate has a mild inhibitory effect on direct activation of SR  $Ca^{2+}$  release by caffeine or  $Ca^{2+}$  (143, 163), skinned fiber experiments with functional EC coupling have shown that voltage-sensor activation of  $Ca^{2+}$  release is little if at all affected, and twitch and tetanic forces are virtually unchanged in the presence of 30 mM cytoplasmic lactate (143, 358). It has also been found recently that increases in intracellular lactate concentration do not cause fiber swelling, owing to compensatory effects of intracellular  $H^+$  changes on the number of osmotically active particles (444). Furthermore, the gradual accumulation of extracellular lactate around muscle fibers during exercise in vivo probably reduces any tendency for intracellular lactate to trigger vacuolation of the

T system (263). In summary, it appears that the intracellular accumulation of lactate per se is not a major factor in muscle fatigue.

## 2. $pH_i$ and muscle fatigue

In humans, muscle  $pH_i$  at rest is  $\sim 7.05$  and after exhaustive exercise may drop to as low as  $\sim 6.5$  (389, 413). In other cases, however,  $pH_i$  decreases only to  $\sim 6.8$  or  $6.9$  at the point of exhaustion (30, 215), showing that muscle fatigue in humans often occurs without there being any large increase in  $[H^+]_i$ . Similarly, in rat gastrocnemius muscle stimulated in vivo, tetanic force decreased by 60%, even though the  $pH_i$  in the muscle observed by NMR declined only to  $\sim 6.9$  (21). Very similar results were also found when monitoring  $pH_i$  with a fluorescent indicator in isolated single fast-twitch mouse fibers subjected to repeated intermittent stimulation (71, 463). Furthermore, humans with myophosphorylase deficiency are unable to utilize muscle glycogen, and their muscles fatigue faster than normal individuals; this accelerated fatigue development occurs without any change in  $pH_i$  (78). Thus, clearly, muscle fatigue can be caused by factors other than raised  $[H^+]$ .

At the onset of exercise or muscle stimulation,  $pH_i$  may initially increase by  $\sim 0.1$  pH units (215, 463) owing to the  $H^+$  consumed during PCr breakdown, and conversely, after exercise ceases,  $pH_i$  can decrease a further  $\sim 0.1$  pH units as PCr is resynthesized (310, 391). Importantly, in cases where  $pH_i$  does drop to low levels in a fatigued muscle, upon ceasing the exercise or stimulation, force typically recovers much faster than  $pH_i$  (22, 78, 391, 438), indicating that the low pH per se was not responsible for all of the force reduction. Furthermore, attenuating the decline in pH during a stimulation period did not reduce the extent of fatigue in frog muscle fibers (421).

A poor correlation between decreased  $pH_i$  and impaired contractile function is also apparent, particularly in mammalian muscle fibers at near physiological temperature, when  $pH_i$  is experimentally decreased, for instance by raising  $CO_2$  concentration. Ranatunga (365) found that twitch and tetanic force in rat EDL bundles at  $30\text{--}35^\circ\text{C}$  were not decreased at acid  $pH_i$  (likely  $\sim 6.5$ ), but actually potentiated. In cat muscle in vivo at  $37^\circ\text{C}$ , decreasing muscle  $pH_i$  to  $6.3$  by hypercapnia reduced maximum tetanic force by only  $\sim 5\text{--}10\%$  in soleus and biceps brachii muscles (2). In intact single fibers from mouse at  $32^\circ\text{C}$ , decreasing  $pH_i$  from  $7.17$  to  $6.67$  caused only  $\sim 10\%$  reduction in maximum tetanic force and no significant slowing of the maximum velocity of shortening (472). In all these cases, decreasing  $pH_i$  to the levels seen in fatigued fibers caused relatively little reduction in tetanic force. Similar conclusions have been reached for frog fibers (20, 373).

## 3. Effects of low pH on SR $Ca^{2+}$ release and voltage sensor activation

Another way in which low pH previously was thought to reduce force responses was by inhibiting  $Ca^{2+}$  release from the SR. Low pH does reduce direct activation of the  $Ca^{2+}$  release channel to stimulation by  $Ca^{2+}$  and caffeine (277, 294, 482); however, voltage-sensor activation of  $Ca^{2+}$  release, the normal physiological mechanism, is not noticeably inhibited even at pH  $6.2$  (256, 258), nor is the activation of the voltage sensors themselves (26). In accord, when  $pH_i$  is lowered in intact fibers, maximum tetanic force is reduced by no more than the amount expected from the reduction in  $F_{Ca,max}$  of the contractile apparatus occurring at that temperature (20, 364, 466, 472).

## 4. Effects of low pH on $Ca^{2+}$ binding to TnC and the SR pump

Low pH also reduces the  $Ca^{2+}$  sensitivity of the contractile apparatus (130, 160) probably because the  $H^+$  competes with  $Ca^{2+}$  binding to troponin C. This is often presumed to have major deleterious effects on muscle performance. However, low pH also reduces the affinity of  $Ca^{2+}$  binding at other sites in the muscle fiber, in particular to the SR  $Ca^{2+}$  pump. Thus, even though the affinity of troponin C (TnC) for  $Ca^{2+}$  may be reduced, the total amount of  $Ca^{2+}$  binding to TnC may not decrease. A single AP normally triggers the release of a comparatively large amount of  $Ca^{2+}$  from the SR ( $\sim 20\%$  of the SR  $Ca^{2+}$  content or  $\sim 230\text{ }\mu\text{M}$   $Ca^{2+}$  expressed relative to total cell volume) (37, 360), and the rise in free  $[Ca^{2+}]$  in the cytoplasm is only a very small fraction of this, because most of the  $Ca^{2+}$  binds to TnC, the  $Ca^{2+}$  pump, and other sites (37, 307, 360). Lowering pH from  $7.0$  to  $6.3$  reduces the affinity of the SR  $Ca^{2+}$  pump more than fivefold ( $0.14\text{--}0.72\text{ }\mu\text{M}$ ) (488), which is considerably greater than the effect on the contractile apparatus ( $<2$ -fold decrease in  $Ca^{2+}$  affinity) (20, 160). Consequently, the amount of  $Ca^{2+}$  available to bind to TnC is likely higher at acid pH. Furthermore, the greatly reduced  $Ca^{2+}$  affinity and rate of uptake by the SR  $Ca^{2+}$  pump at acid pH (8, 488) leads to an increase in the resting cytoplasmic free  $[Ca^{2+}]$  (466), which would help maintain or even increase the extent of resting occupancy of the various cytoplasmic  $Ca^{2+}$  binding sites. As a result of all these effects, the free  $[Ca^{2+}]$  in the cytoplasm during contraction actually reaches greater levels at acid pH than at normal pH (20, 466), and the force response to a twitch is actually increased (20, 365) and tetanic force rises faster, at least in frog muscle (364).

In summary, under physiological circumstances, low pH has far less inhibitory effects on the activation of the contractile apparatus and  $Ca^{2+}$  release than previously assumed, and its effects on the SR  $Ca^{2+}$  pump actually favor force development.

### 5. *Effect of low pH on membrane conductance*

Importantly, as mentioned in section viC7, reduced  $pH_i$  can have marked beneficial effects on muscle performance by helping maintain excitability. Nielsen and colleagues (200, 333) demonstrated that lowering  $pH_i$  to  $\sim 6.6$  in both fast-twitch and slow-twitch mammalian muscle helped counteract the reduction in excitability occurring with membrane depolarization. This occurred because lowering  $pH_i$  reduced the  $Cl^-$  conductance of the T-system membrane by approximately twofold, enabling APs to continue to propagate over both the surface and T-system membranes and hence trigger contraction (351, 352). Juel and colleagues (248) confirmed the finding that lowering  $pH_i$ , by addition of extracellular lactic acid, helped restore excitability in depolarized mammalian muscle, but they questioned whether this occurs in active muscle because they found that the rate of force decrease during fatigue was unaffected in muscles stimulated in vitro under such conditions. However, in those experiments, the fibers deeper in the muscles would have become anoxic due to the high stimulation rate and limited  $O_2$  diffusion (32), which would have greatly increased the rate of fatigue (493), thereby masking any beneficial effects of low  $pH_i$ . Furthermore, it is quite likely that decreasing  $pH_i$  over and above that normally occurring during fatigue has no extra beneficial effects on the excitability. Experiments in which lactate was infused into whole animals while the muscles were stimulated to fatigue by nerve or direct muscle stimulation, found that the presence of extracellular lactate, which likely decreased  $pH_i$ , reduced failure of sarcolemma excitability and restored force production (239), consistent with an action on  $Cl^-$  conductance.

### 6. *Effects of low pH on the rate of fatigue*

It has been shown in isolated muscle fibers that lowering  $pH_i$  from 7.18 to 6.77 did not decrease either the initial level of tetanic force or the rate at which the muscle fiber fatigued when given repeated stimulation, even though the final  $pH_i$  was much lower at the point where force dropped to 40% ( $pH_i$  6.46 versus 6.91 in control case) (71). In fact, it took  $98 \pm 18$  tetani to cause force to drop to 40% in the acidic conditions compared with  $63 \pm 10$  tetani in the control conditions, suggesting that the rate of fatigue if anything was less at the lower  $pH_i$ . This effect might in part have been due to improved excitability in the T system (352) or perhaps was mainly due to a pH-dependent decline in the rate of ATP consumption owing to reduced cross-bridge cycling (71). The results also offered no evidence that low  $pH_i$  accelerates muscle fatigue by decreasing glycolysis. Similarly, in humans, glycolysis and glycogenolysis are not inhibited by the  $pH_i$  decreases occurring in exhaustive exercise (29, 30, 194), and studies with electrical stimulation in humans have

shown that even though glycolysis and glycogenolysis are reduced somewhat when  $pH_i$  falls from 6.7 to 6.45, significant activity still remains (415).

Finally, there have been many studies examining the effects of manipulating whole body pH on exercise performance, with some albeit not all showing deleterious effects of acidity and beneficial effects of alkalosis (79). However, altering body pH can have effects on blood oxygen saturation and oxygen unloading, cardiac and local vasculature function, central nervous system drive, and other factors. Experiments with a perfused rat hindlimb preparation, where the effects of some of these variables could be controlled, showed that a decrease in blood pH adversely affected muscle performance, but that this was due to some direct effect of the low extracellular pH itself and not of lowered  $pH_i$  (414). The accompanying perfusion pressure data indicated that the effect was quite possibly due entirely to the acidity disrupting the normal local control of blood flow in the muscle vascular beds. Similarly, alkalosis in humans can delay the onset of fatigue (409), but such alkalosis actually causes a decrease in extracellular  $[K^+]$ , and this may have additional effects in aiding membrane excitability. Significantly, when such changes in  $[K^+]$  and other effects were avoided in a perfused hindlimb preparation, alkalosis did not improve muscle performance (412). Finally, increasing muscle carnosine by  $\beta$ -alanine supplementation has been found to delay the onset of fatigue in exercising humans, and this was attributed to increased pH buffering by the carnosine (209), but the beneficial effects of the carnosine are quite likely mainly due to it markedly increasing the  $Ca^{2+}$  sensitivity of the contractile apparatus (144, 261).

In conclusion, raised  $[H^+]_i$  per se is not the main cause of muscle fatigue, with its direct effects on force production being quite small.

### C. ATP and $Mg^{2+}$

The changes in ATP, ADP, IMP, and  $[Mg^{2+}]_i$  that occur when consumption of ATP exceeds resynthesis have been described in section III. To summarize, during intense fatigue, taking account of the greater changes in fast fibers (238), ATP may decline from 7 to 1.2 mM and PCr from 30 to 2.5 mM while ADP may increase from 10 to 200  $\mu$ M, IMP may increase from undetectable to 5 mM, and  $[Mg^{2+}]_i$  may increase from 1 to 2 mM or possibly higher. In slow fibers, on the other hand, there is probably little if any decline in cytoplasmic [ATP] (238, 324). As noted in section III, it is possible that in fast fibers the changes in some restricted regions of the cell are even larger than the measured average cytoplasmic changes. In this section we consider the resulting changes in contractile apparatus and EC coupling that may contribute to fatigue.



### 1. Contractile apparatus

Maximum force production in skinned fibers is not reduced unless the [ATP] is  $<20 \mu\text{M}$  (98), and is actually increased by  $\sim 10\%$  at  $0.5\text{--}1 \text{ mM}$  ATP (145, 185) and also by submillimolar ADP and AMP (185).  $\text{Ca}^{2+}$  sensitivity is not altered by lowering the [ATP] to  $0.5 \text{ mM}$ , but is slightly increased by submillimolar ADP (145, 185, 296). One early suggestion was that IMP accumulation may cause fatigue (168); however,  $3 \text{ mM}$  IMP has no effect on maximum force and causes a slight increase in  $\text{Ca}^{2+}$  sensitivity (56). Increasing  $[\text{Mg}^{2+}]$  to  $3 \text{ mM}$  has no effect on maximum force or the velocity of shortening (145) but causes a substantial decrease in  $\text{Ca}^{2+}$  sensitivity (1.6-fold increase in  $[\text{Ca}^{2+}]$  required for 50% maximal force) (56). In fast fibers, the combined effect of all these changes is likely to be a slight increase in maximum force but a substantial decrease in the  $\text{Ca}^{2+}$  sensitivity of force production. These effects are likely additional to those occurring with raised  $\text{P}_i$  (see sect. VIIA) and  $\text{H}^+$  (see sect. VIIB).

### 2. SR $\text{Ca}^{2+}$ pumping and leakage

Studies on skinned fibers show that the rate of relaxation of tetani is reduced 2.5-fold when [ATP] is lowered to  $0.5 \text{ mM}$  (at constant low [ADP]) (144). This is not due to an effect on the contractile apparatus, but instead is likely due to reduced  $\text{Ca}^{2+}$  uptake by the SR  $\text{Ca}^{2+}$  pumps. Although the ATP catalytic site on the SR pumps has a very high affinity ( $K_m \sim 1\text{--}10 \mu\text{M}$ ), ATP also has a regulatory action on the pump, such that if [ATP] decreases from  $5$  to  $0.25 \text{ mM}$ , the  $\text{Ca}^{2+}$  affinity of the pump is reduced  $\sim 10$ -fold together with a drop in the  $\text{Ca}^{2+}$  binding cooperativity (327, 432). The latter change may enable the ATPase to continue pumping  $\text{Ca}^{2+}$  despite a considerable decrease in the free energy of ATP hydrolysis, as only one  $\text{Ca}^{2+}$  is transported for each ATP molecule hydrolyzed, but this means a decrease in the energy efficiency of the process. If the  $\text{Ca}^{2+}$  affinity of the SR pump is already greatly reduced by acid pH (e.g., pH  $\sim 6.3$ ) (488), reducing [ATP] has little if any additional effect (327). In fast-twitch mammalian fibers, a rise in [ADP] from  $10 \mu\text{M}$  to  $0.2 \text{ mM}$  causes a small reduction in  $\text{Ca}^{2+}$  pump rate and induces a twofold increase in leakage of  $\text{Ca}^{2+}$  from the SR via the pump "slippage" (295). Such changes would slow  $\text{Ca}^{2+}$  uptake and increase resting  $[\text{Ca}^{2+}]$  in an intact fiber. The increased SR  $\text{Ca}^{2+}$  leakage is observed even in the presence of  $5\text{--}8 \text{ mM}$  ATP if CrP is decreased to low levels (132, 134, 295), indicating that the local [ADP] near the  $\text{Ca}^{2+}$  pumps is poorly controlled if CrP is depleted. Interestingly, in slow-twitch fibers, raised [ADP] has only comparatively small effects on the SR pump rate and leakage (296). Finally,  $\text{Ca}^{2+}$  uptake in fast-twitch fibers is little if at all affected by raising  $[\text{Mg}^{2+}]$  from  $1$  to  $3 \text{ mM}$  (237) or by the presence of  $3 \text{ mM}$  IMP (55).

In summary, studies on skinned muscle fibers suggest that the decline in [ATP] and [CrP] can reduce SR  $\text{Ca}^{2+}$  pumping and increase pump leakage, resulting in elevated resting  $[\text{Ca}^{2+}]_i$  as typically observed in fatigue. Under some conditions, it may also contribute to the slowing of relaxation in fatigued muscle (see sect. XI).

### 3. SR $\text{Ca}^{2+}$ release

The  $\text{Ca}^{2+}$  release channel in mammalian skeletal muscle (RyR1) is strongly stimulated by ATP binding at a cytoplasmic regulatory site ( $K_{\text{app}} \sim 0.5 \text{ mM}$ ), and ADP and AMP act as weak competitive agonists, but IMP does not (278, 305). The  $\text{Ca}^{2+}$  release channel is also strongly inhibited by cytoplasmic  $\text{Mg}^{2+}$  (257, 258, 280, 281, 305) and remains closed at the resting  $[\text{Mg}^{2+}]$  ( $\sim 1 \text{ mM}$ ) unless directly activated by the voltage sensors. Voltage sensor-induced  $\text{Ca}^{2+}$  release is decreased  $\sim 40\%$  by raising  $[\text{Mg}^{2+}]$  from  $1$  to  $3 \text{ mM}$  (145). Voltage sensor-induced  $\text{Ca}^{2+}$  release is also decreased  $\sim 20\%$  by lowering [ATP] to  $0.5 \text{ mM}$ , and the reduction in  $\text{Ca}^{2+}$  release is larger when raised  $[\text{Mg}^{2+}]$  and low [ATP] are combined, and larger still if ATP hydrolysis products, AMP or adenosine, are also present (55, 56, 145, 342). Consequently, it seems likely that this combination of factors could substantially inhibit voltage sensor-induced  $\text{Ca}^{2+}$  release. The inhibitory effects of increased  $[\text{Mg}^{2+}]$  and reduced [ATP] on  $\text{Ca}^{2+}$  release likely underlie the decrease in tetanic  $[\text{Ca}^{2+}]_i$  observed in  $\text{CK}^{-/-}$  mice at the onset of high-intensity stimulation (107), but it is apparent that they are not the only metabolic factors involved in reducing  $\text{Ca}^{2+}$  release in fatigue (108) (see sects. VIIB and VID).

In conclusion, in intense exercise, the triad junction may play a key role by sensing depletion of cellular [ATP] levels and respond by reducing  $\text{Ca}^{2+}$  release. This will decrease the rate of ATP usage by reducing both cross-bridge cycling and SR  $\text{Ca}^{2+}$  uptake, the two main sources of ATP hydrolysis. The cost of this is a reduction in power output, or in other words muscle fatigue, but the benefit is that it ultimately prevents complete exhaustion of all cellular ATP and consequent rigor development and cellular damage.

## D. Glycogen

In skeletal muscle, glucose is stored as glycogen, and this store is a major source of energy during most forms of muscle activity. A direct correlation between muscle glycogen concentration and time to fatigue during moderately intense exercise ( $60\text{--}80\%$  of maximal oxygen uptake) was first shown by Hultman and co-workers (46, 204). Their findings have subsequently been confirmed in numerous studies, and the association between glycogen depletion and fatigue during moderately intense exercise is now well established. However, the link between gly-



cogen depletion and the force decrease during fatigue is not fully understood.

Experiments with simultaneous measurements of force and  $[Ca^{2+}]_i$  were performed to shed light on cellular mechanisms linking low glycogen to fatigue. In an initial study, small bundles of fast-twitch flexor digitorum brevis (FDB) muscle fibers were fatigued by repeated tetani, which reduced their glycogen content to  $\sim 25\%$  of the control (90). When recovery occurred in the absence of glucose, glycogen did not recover, and fiber bundles fatigued more rapidly in a subsequent fatigue run. Accompanying experiments on single muscle fibers showed that the decrease in tetanic force during fatigue coincided with reduced  $Ca^{2+}$  transients both in the initial fatigue run and in the second run, which occurred much faster when recovery took place in the absence of glucose (90). Similar results were subsequently obtained in single cane toad muscle fibers that were repeatedly fatigued by intermittent tetanic stimulation in the absence of glucose (236). Moreover, isolated mouse fast-twitch EDL muscles were fatigued by repeated tetani; allowed to recover for 2 h in zero, normal, or high extracellular glucose; and then fatigued again. Muscles recovering in zero glucose had lower glycogen levels ( $\sim 50\%$  of the control) at the start of the second fatigue run and fatigued more rapidly than muscles recovering in normal or high glucose. The accelerated fatigue affected both tetanic force and  $[Ca^{2+}]_i$  (203). Thus a reduced prefatigue level of glycogen is associated with a faster decrease of tetanic  $[Ca^{2+}]_i$  and force during fatigue.

The link between reduced glycogen and decreased  $[Ca^{2+}]_i$  transients may be due to glycogen providing acetyl coenzyme A for the tricarboxylic acid (TCA) cycle or maintaining high levels of TCA intermediates (203, 390). Alternatively, the association between low glycogen and impaired SR  $Ca^{2+}$  release is not directly linked to glycogen's role in energy metabolism. This alternative originates from a study on skinned toad muscle fibers where the ability to respond to t-tubular depolarizations correlated closely with the muscle glycogen content (non-soluble component) (425). In these skinned fiber experiments, ATP and PCr were present in the bathing solutions, suggesting that glycogen had a structural rather than a metabolic role. Similar studies were subsequently performed on skinned rat EDL fibers, and these mammalian fibers showed only a small (34) or no (191) ATP- and PCr-independent effect of glycogen on the capacity to respond to depolarizations. Data from fatigue studies on intact muscle cells may be used in support of both a metabolic and structural role of glycogen. Isolated FDB fibers and EDL muscles fatigued more rapidly when glycogen was decreased after recovery in zero glucose, which is consistent with both possibilities (90, 203). However, the glycogen-depleted cells displayed normal changes in other fatigue-induced parameters (i.e., de-

creased myofibrillar  $Ca^{2+}$  sensitivity and maximum force, slowed relaxation, and increased resting  $[Ca^{2+}]_i$ ), which are generally attributed to metabolic changes (90, 203). On the other hand, fatigue in cane toad muscle fibers produced under control conditions was accompanied by a decrease in the rapidly releasable SR  $Ca^{2+}$  stores, whereas this store was not decreased after the accelerated fatigue development in glycogen-depleted cells (236). Thus these data on toad fibers indicate different mechanisms behind the fatigue-induced reduction in SR  $Ca^{2+}$  release in normal and in glycogen-depleted cells.

In conclusion, depletion of glycogen during prolonged, exhausting exercise may contribute to fatigue by causing decreased SR  $Ca^{2+}$  release. The mechanistic link between low glycogen and premature failure of SR  $Ca^{2+}$  release remains uncertain.

## VIII. REACTIVE OXYGEN SPECIES

There is a large and rapidly growing literature establishing that reactive oxygen species (ROS) are produced in active muscles and have some role in fatigue. The most convincing evidence that ROS contribute to fatigue comes from experiments with exogenously added ROS scavengers which can reduce the rate of fatigue in isolated muscles, in intact animals, and in humans. Despite the overwhelming evidence for some role for ROS in muscle fatigue, many aspects remain uncertain, particularly the source of the ROS, the identification of the particular ROS which are important, and the mechanism(s) by which ROS contribute to fatigue. There are many valuable reviews of ROS in muscle (94, 368, 428), and here we only briefly reiterate the areas covered in those reviews. A comprehensive account of the chemistry and biology of ROS is available (196). Our main focus is current ideas on the mechanisms by which ROS contribute to fatigue.

The most important ROS are superoxide ( $O_2^{\bullet -}$ ), hydrogen peroxide ( $H_2O_2$ ), and hydroxyl radicals ( $OH^{\bullet}$ ). In addition, nitric oxide can interact with superoxide to form peroxynitrite ( $ONOO^-$ ), one of a series of reactive nitrogen species, which may also have normal and pathological roles in muscle (369). Superoxide is produced in mitochondria as a by-product of oxidative phosphorylation (443) and also by various enzymes including NADPH oxidase, xanthine oxidase, and lipoxygenases (Fig. 4). Superoxide is only moderately reactive and is rapidly broken down by various superoxide dismutases (SOD) to hydrogen peroxide. Given its negative charge, superoxide does not easily cross membranes but may utilize anion channels in the surface membrane (292) or voltage-dependent anion channels (VDACs) in the mitochondrial membrane for this purpose (443).

Hydrogen peroxide is not highly reactive but diffuses through cell membranes and is thought to have a signaling




FIG. 4. The major reactive oxygen species in muscle. Numbers in brackets indicate approximate lifetimes of various species (39, 196). GPX, glutathione peroxidase; GR, glutathione reductase. Other abbreviations are as in text.

role in addition to oxidizing some thiol groups on proteins (196). Hydrogen peroxide can be broken down by various pathways; of particular importance is the reaction with free transition metals such as  $\text{Fe}^{2+}$  (Fenton reaction), which produces the extremely reactive hydroxyl radical. Hydroxyl radicals have exceedingly short lifetimes because they react with virtually any organic molecule and can damage proteins, DNA, and lipids. The  $\text{Fe}^{3+}$  present in myoglobin, hemoglobin, or cytochrome *c* can be reduced to  $\text{Fe}^{2+}$  by superoxide radicals (Haber-Weiss reaction) and then allows hydroxyl radical production. Hydrogen peroxide is also broken down harmlessly by catalase ( $2\text{H}_2\text{O}_2 \rightarrow 2\text{H}_2\text{O} + \text{O}_2$ ) or by glutathione peroxidase with the conversion of reduced glutathione (GSH) to oxidized glutathione (GSSG). Given that the important ROS form a cascade from superoxide to hydrogen peroxide to hydroxyl radicals, it is difficult to pinpoint which particular species is exerting an effect.

#### A. ROS Production Is Accelerated in Active Muscles

It is generally agreed that ROS production is accelerated by muscle activity and by increased temperature which, of course, usually accompanies activity. In early studies, increased ROS production in active muscle was established by electron resonance spectroscopy combined with spin trapping and by nonspecific markers of oxidation, such as lipid peroxidation products and protein oxidation products (115, 127).

Better time resolution and potentially distinction between intracellular and extracellular ROS can be obtained by the use of marker molecules that are located in either the intracellular or extracellular space. Reid et al. (371) used intracellular dichlorofluorescein, which fluoresces in response to various ROS, and found a modest increase in ROS during repetitive tetani. Interestingly, application of SOD and catalase, which would remain extracellular, substantially reduced fluorescence that was interpreted as reducing the extracellular concentrations and thereby

preventing back diffusion of superoxide and/or  $\text{H}_2\text{O}_2$ . Extracellular ROS production can be determined by perfusion with cytochrome *c*, which changes its absorption after reduction by superoxide. Increased extracellular superoxide production during activity has been reported (245, 371, 427), which suggests that superoxide is either leaving the cell through anion channels or that it is produced from an extracellular site.

Increases in intracellular and extracellular ROS production as a function of temperature have also been reported by several groups (17, 447, 497). Zuo et al. (498) detected extracellular superoxide and tested whether it was leaving the cell through anion channels by use of the blockers probenecid and DIDS. Neither blocker affected extracellular superoxide, suggesting an extracellular source.

#### B. Sources of ROS in Muscle

While increased production of ROS during muscle activity is generally accepted, the source of the ROS remains unclear. Mitochondria are generally thought to produce superoxide at a rate of  $\sim 1\text{--}2\%$  of  $\text{O}_2$  consumption, although markedly lower estimates ( $\sim 0.15\%$ ) have also been obtained (418). They contain abundant MnSOD, so much of this superoxide may be broken down to  $\text{H}_2\text{O}_2$ , at least under resting conditions (443). During intense activity, when mitochondrial  $\text{O}_2$  consumption increases up to 100-fold (94), it is postulated that the excess superoxide may escape the mitochondria via VDACS or be converted to  $\text{H}_2\text{O}_2$  that can diffuse across the mitochondrial membrane into the myoplasm (94, 115). Early attempts to test this hypothesis by inhibiting mitochondrial complexes with rotenone failed to reduce muscle ROS production (226, 498). However, a recent study using skinned fibers in which the mitochondria were inhibited with rotenone and succinate (to prevent reverse electron transport across complex I) found that superoxide production was reduced to zero (446).

Mitochondrial ROS production may also be stimulated indirectly by products of phospholipase A<sub>2</sub> (PLA<sub>2</sub>), since inhibition of PLA<sub>2</sub> reduced ROS production (187, 329). Another possible source of ROS is nonphagocytic NAD(P)H oxidase (NOX). There are now several reports of NOX in skeletal muscle that may be associated with the SR (492) or with the t-tubular membrane (207, 226). These studies all showed NAD(P)H-dependent superoxide production, and in two of the studies, inhibitors of NOX (diphenyleneiodonium, DPI) reduced superoxide production by muscle (207, 226). Xanthine oxidase is a further possible source of ROS, and the inhibitor allopurinol has inhibited ROS production in some (427) but not other studies (226). Lipoxygenases (496) are another possible source of superoxide.

Currently there is no consensus about the major sources of ROS production within muscle, but mitochondria and NAD(P)H oxidase seem probable contenders. It is likely that different muscles under various conditions utilize different pathways for ROS production, and it is also likely that the intracellular and extracellular sources of ROS are different.

### C. ROS Scavengers Reduce Fatigue

The earliest study of ROS effects on fatigue utilized strips of diaphragm muscle perfused by the circulation and stimulated through the phrenic nerve (400). *N*-acetylcysteine (NAC) injected into the circulation (150 mg/kg body wt) had no effect on the control force but increased the force at the end of a period of intermittent 20-Hz stimulation from 45% control to 66%. Improvement by NAC was present both at low frequencies of stimulation and at high, but NAC did not appear to improve recovery from fatigue. A range of ROS scavengers have subsequently been shown to slow fatigue including SOD, DMSO, catalase (371), Tiron, and Tempol (315). The magnitude of the improvement by ROS scavengers in different studies is quite variable. Two factors that contribute to this are frequency of stimulation (370) and temperature (315). Reid et al. (370) fatigued human tibialis anterior muscles by electrical stimulation using intermittent tetani at either 40 or 10 Hz. During tetani at 40 Hz, the muscles fatigued rapidly, but NAC (150 mg/kg infused intravenously over 1 h) had no effect. In contrast, when tetani were only partially activated by 10-Hz stimulation, fatigue was much slower and NAC produced a moderate improvement. MVCs were also unaffected by NAC. The authors propose that ROS affect either SR Ca<sup>2+</sup> release or myofibrillar Ca<sup>2+</sup> sensitivity so that only low-frequency tetani are sensitive.

Muscle temperature appears to be another factor that can explain some of the variability. In a study on mouse single fibers, Moopanar and Allen (315) found that fatigue

at 22°C was unaffected by ROS scavengers, while fatigue at 37°C was substantially slowed by ROS scavengers. The increased sensitivity of muscle fatigue to ROS at 37°C may result from the increased rate of ROS production as temperature increases. Since increased activity and temperature both increase ROS production, it is clear that regular exercise will be associated with substantial increases in ROS production. Given the potentially damaging effects of ROS, it is perhaps not surprising that endogenous ROS scavenging pathways are substantially up-regulated by regular exercise. For instance, SOD activity and GPX activity are increased in the actively recruited muscles and preferentially in oxidative muscles. There is also evidence that the levels of GSH can increase. These issues have been extensively reviewed elsewhere (362). Furthermore, a recent study showed that PPAR $\gamma$  coactivator 1 $\alpha$ , which is a potent stimulator of mitochondrial biogenesis, has a central role in the induction of several ROS defense enzymes in response to oxidative stress (419).

Although improvements in fatigue in response to ROS scavengers have been frequently reported in isolated muscles or intact muscles in animals, improvements in human performance in response to dietary antioxidants (e.g., vitamins C and E) have not usually been observed. An exception is the thiol donor NAC, which seems to have beneficial effects on fatigue performance as already described (370, 400). An important study on trained humans extends the beneficial effects of NAC to a voluntary submaximal cycling in which the time to volitional fatigue was increased by NAC infusion by 26% (304). In a recent extension of this approach, McKenna et al. (303) showed that NAC minimized the reduction in Na<sup>+</sup>-K<sup>+</sup> pump activity that usually occurs in exercise (394) and also attenuated the rise in plasma K<sup>+</sup>. Given that the Na<sup>+</sup>-K<sup>+</sup> pump is redox sensitive (184), the authors propose that ionic changes associated with Na<sup>+</sup>-K<sup>+</sup> pump contribute to fatigue and can be ameliorated by the ROS scavenger NAC.

### D. Mechanisms by Which ROS Cause Fatigue

As discussed above, there is very strong evidence that ROS contribute to some models of fatigue, but understanding of the mechanisms involved is currently limited. Below we discuss experimental investigation of the mechanisms under three headings, recognizing, of course, that the eventual aim will be to identify specific proteins and specific biochemical changes that underlie these effects.

#### 1. Maximum Ca<sup>2+</sup>-activated force

Studies of the effects of ROS on F<sub>Ca,max</sub> have been undertaken both in intact muscles and in skinned fibers.



In intact preparations, short exposures to  $\text{H}_2\text{O}_2$  can increase force moderately while longer exposures can decrease force, and at least some of these effects seem to be through changes in  $F_{\text{Ca,max}}$  (14, 356). In skinned fiber studies, it is clear that very high concentrations of sulfhydryl oxidizing agents, such as 10 mM dithionitrobenzoic acid (485) or 1 mM dithiodipyridine (255), can reduce  $F_{\text{Ca,max}}$  to zero, but these concentrations are not physiological. Moderate concentrations of ROS have given quite variable results. For instance, in skinned rabbit muscle, Darnley et al. (113) showed that both superoxide and  $\text{H}_2\text{O}_2$  could produce a 50% enhancement of  $F_{\text{Ca,max}}$  during an acute exposure, but after the ROS were removed, there was a substantial reduction in force. However, in the same preparation, Callahan et al. (84) found no effects of  $\text{H}_2\text{O}_2$ , while superoxide caused a 15% reduction in  $F_{\text{Ca,max}}$  and hydroxyl radicals reduced force by 44%. Lamb and Posterino (255), however, found that in rat skinned fibers, with the exception of the high concentration of oxidizing agent mentioned above, none of the oxidants or reductants used affected  $F_{\text{Ca,max}}$ .

There have also been studies that try to dissect the role of ROS during fatigue or high temperature. Moopanar and Allen (315, 316) tested  $F_{\text{Ca,max}}$  in mouse single fibers fatigued by intermittent tetani. At 22°C, fibers fatigued slowly, and fatigue in this model has been shown to depend on declines in  $F_{\text{Ca,max}}$ ,  $\text{Ca}^{2+}$  sensitivity, and tetanic  $[\text{Ca}^{2+}]_i$  due to mechanisms discussed elsewhere in this review. When the same fibers were stimulated at 37°C, they fatigued much more quickly, and this increased fatigability could be reversed by the membrane-permeant antioxidants Tiron or Tempol. However,  $F_{\text{Ca,max}}$  declined to the same extent at 22 and 37°C, establishing that a decline in  $F_{\text{Ca,max}}$  was not the cause of the ROS-sensitive component of fatigue at 37°C. This finding is consistent with several reports in which isolated muscles were fatigued and then skinned, but no persisting changes in  $F_{\text{Ca,max}}$  were noted (111, 483, 484).

During intense exercise, high muscle temperatures can occur, and Van der Poel and Stephenson (447) explored the effects of increasing temperature to 43–47°C for short periods in unstimulated muscles (these temperatures are well above those experienced by muscles during normal exercise). The fibers were then skinned and  $F_{\text{Ca,max}}$  and  $\text{Ca}^{2+}$  sensitivity determined as functions of time. The main result was that  $F_{\text{Ca,max}}$  showed a large fall but then spontaneously recovered. Importantly, if the muscle was treated with Tiron, it prevented both the superoxide production and the reduction in  $F_{\text{Ca,max}}$ . This result, which contrasts with Moopanar and Allen (315) above, suggests that high temperatures in unstimulated muscle cause force decline by a different mechanism to fatigue at 37°C.

## 2. $\text{Ca}^{2+}$ sensitivity

In the intact single fiber study of fatigue at 37°C, Moopanar and Allen (315) found that the rapid decline of force was largely caused by a fall in  $\text{Ca}^{2+}$  sensitivity. This fall in sensitivity could be prevented by preexposure to Tiron or Tempol and could be reversed by brief application of dithiothreitol (DTT) (316). Studies of skinned fibers have generally not noted pronounced changes in  $\text{Ca}^{2+}$  sensitivity when ROS were applied to the skinned fiber (84, 113). An exception is the study by Posterino and Lamb (255) who found that prolonged exposure to ROS ( $\text{H}_2\text{O}_2$  or dithiodipyridine, DTDP) during cycles of activation generally increased  $\text{Ca}^{2+}$  sensitivity. In addition, exposure to GSH could lead to both increases and decreases in  $\text{Ca}^{2+}$  sensitivity under different conditions. These results show 1) that the presence of GSH can modify the effects of oxidants on sensitivity changes, 2) that probably there are several sites which affect  $\text{Ca}^{2+}$  sensitivity and are modulated by oxidation state, and 3) that both the magnitude and direction of sensitivity change can depend on muscle history and concentration and exposure time of oxidants. Because most studies on skinned fibers have not included GSH, this may explain some of the negative results.

Studies of the effects of exogenous  $\text{H}_2\text{O}_2$  on intact fibers also showed changes in  $\text{Ca}^{2+}$  sensitivity (14, 15). Initially  $\text{Ca}^{2+}$  sensitivity increased, but longer exposures to larger concentrations of  $\text{H}_2\text{O}_2$  caused a large fall in  $\text{Ca}^{2+}$  sensitivity that was reversed by DTT. Interestingly, DTT caused a small decrease in  $\text{Ca}^{2+}$  sensitivity when applied alone, suggesting that the endogenous oxidative state of proteins may affect  $\text{Ca}^{2+}$  sensitivity perhaps by a ROS-sensitive signaling pathway (15).

A series of studies from Nosek's laboratory have explored the effects of hypoxic fatigue on the properties of muscle when subsequently skinned. Fatigue alone had no effect on the function of the subsequently skinned preparation, whereas hypoxic fatigue caused reductions in both  $F_{\text{Ca,max}}$  and  $\text{Ca}^{2+}$  sensitivity (63). Subsequently, these changes in the skinned preparations were shown to be associated with degradation of troponins I and C, which the authors speculate may be caused by increased levels of ROS (119).

## 3. SR $\text{Ca}^{2+}$ release

Isolated SR  $\text{Ca}^{2+}$  channels in a lipid bilayer show an increased open probability in response to  $\text{Ca}^{2+}$  when  $\text{H}_2\text{O}_2$  is present at submillimolar concentration (162). In addition, it has long been known that ROS can inhibit the SR  $\text{Ca}^{2+}$ -ATPase and reduce the rate of  $\text{Ca}^{2+}$  uptake into the SR (395). Despite these suggestive results on simplified preparations,  $\text{H}_2\text{O}_2$  had remarkably little effect on the amplitude of the tetanic  $[\text{Ca}^{2+}]_i$  in single fibers at concentrations and durations which caused substantial changes



in force due to  $\text{Ca}^{2+}$  sensitivity changes (14, 15). The only exception was that long exposures to high concentrations of  $\text{H}_2\text{O}_2$  cause a gradual increase in the magnitude of the  $\text{Ca}^{2+}$  transient coupled to a slowing of the rate of decline of the  $\text{Ca}^{2+}$  transient, suggesting that the SR  $\text{Ca}^{2+}$  pump was being inhibited. Furthermore, in studies of fatigue at  $37^\circ\text{C}$ , the rapid decline of force that was ROS sensitive was not accompanied by changes in tetanic  $[\text{Ca}^{2+}]_i$  (315). Interestingly, when ROS were scavenged and fatigue occurred more slowly at  $37^\circ\text{C}$ , the normal pattern of fatigue-induced decline in tetanic  $[\text{Ca}^{2+}]_i$  was present (see sect. VII).

In skinned rat fibers with intact t-tubule/SR coupling, Brotto and Nosek (65) reported that 5-min exposure to 1 mM  $\text{H}_2\text{O}_2$  produced a large reduction in the depolarization-induced  $\text{Ca}^{2+}$  release. These issues were reinvestigated by Posterino et al. (357) using both the same preparation and a skinned preparation in which the sealed t-tubules were electrically stimulated so that the effects of an AP-generated  $\text{Ca}^{2+}$  release could be examined.  $\text{H}_2\text{O}_2$  was found to enhance caffeine-induced  $\text{Ca}^{2+}$  release and  $\text{Ca}^{2+}$ -induced  $\text{Ca}^{2+}$  release and to inhibit voltage-induced  $\text{Ca}^{2+}$  release by an effect on the voltage sensor (since low  $\text{Mg}^{2+}$  could still trigger  $\text{Ca}^{2+}$  release). Critically, the AP-induced  $\text{Ca}^{2+}$  release was unaffected by moderate exposure to  $\text{H}_2\text{O}_2$  or changes in the GSH/GSSG ratio or DTT. These skinned fiber data suggest that the AP, acting through the voltage sensor, normally completely opens the SR release channels and renders them insensitive to changes in the sensitivity of voltage sensor/RyR coupling. Thus data from intact fibers fatigued at  $37^\circ\text{C}$ , intact fibers exposed to  $\text{H}_2\text{O}_2$ , and skinned fibers suggest that changes in SR  $\text{Ca}^{2+}$  release are not likely to contribute the ROS-sensitive component of fatigue.

In summary, it is clear that ROS are generated in active muscles and contribute to the process of fatigue. How ROS are produced and which ROS are important remain uncertain. Many mechanisms of action are possible given the susceptibility of proteins to oxidative damage, but current evidence points at the contractile proteins and the  $\text{Na}^+\text{-K}^+$  pump as the sites showing the greatest susceptibility to ROS under physiological conditions.

## IX. $\text{Ca}^{2+}$ COMPARTMENTS AND MOVEMENTS: ROLE IN FATIGUE

### A. SR $\text{Ca}^{2+}$ Content and $\text{Ca}^{2+}$ Release in Fast- and Slow-Twitch Fibers

The amount of  $\text{Ca}^{2+}$  contained and released within a muscle fiber is of major importance to its function. In rested muscle, the total  $\text{Ca}^{2+}$  content of both fast- and slow-twitch rat fibers is  $\sim 1.1$  mmol/kg wet wt, with most

( $\sim 1.0$  mmol/kg) contained within the SR, equivalent to  $\sim 11$  and  $21$  mM when expressed relative to the respective SR volume (176, 183, 343). Total calcium content in human muscle fibers is slightly lower ( $\sim 0.8$  mmol/kg) (338, 393). Most  $\text{Ca}^{2+}$  in the SR is bound to calsequestrin (38, 45), and the free  $[\text{Ca}^{2+}]$  is  $\sim 0.3\text{--}1$  mM (175, 274, 275, 382). The SR is normally loaded at only  $\sim 25\%$  of maximum capacity in fast-twitch fibers, but at  $\sim 70\%$  or greater in slow-twitch fibers (176, 183, 440). The relatively low fractional SR load level in fast-twitch fibers (and consequent low free  $[\text{Ca}^{2+}]$ ) likely aids  $\text{Ca}^{2+}$  uptake in adverse metabolic conditions. The major  $\text{Ca}^{2+}$ -pump isoform in fast-twitch fibers is SERCA1a and in slow-twitch fibers is SERCA2a, with the faster  $\text{Ca}^{2+}$  uptake in fast-twitch fibers due primarily to the approximately sixfold greater number of pumps (282, 293, 491).

The rate and amount of SR  $\text{Ca}^{2+}$  release per AP is approximately three times higher in fast-twitch fibers than in slow-twitch fibers (37). This difference is due primarily to the higher density of both DHPR/voltage sensors and  $\text{Ca}^{2+}$  release channels in fast-twitch fibers (125, 252, 260, 299). The rapid contraction of fast-twitch fibers requires the presence both of fast MHC (type II) isoforms and fast  $\text{Ca}^{2+}$  release. Compared with fast-twitch fibers, slow-twitch fibers have fewer  $\text{Ca}^{2+}$  binding sites on troponin C and the SR pumps, and a lower rate and amount of  $\text{Ca}^{2+}$  release suffices for contraction, particularly given the much slower contraction rate of the predominant MHC isoform (type I) present (59, 61). Thus SR and contractile properties in a given fiber are generally well matched (440).

### B. Effects of SR Content Changes on $\text{Ca}^{2+}$ Release

Importantly, if the total amount of  $\text{Ca}^{2+}$  in the SR of a fast-twitch fiber drops to substantially below its normal level, the amount released by each AP is reduced (360) (see Fig. 5), reducing the force response. Specifically, a 35% decrease in SR  $\text{Ca}^{2+}$  content (from  $\sim 1.1$  to  $0.71$  mmol/kg) reduced peak tetanic force by 46% (142), despite the fact that the amount of  $\text{Ca}^{2+}$  still available in the SR considerably exceeded the number of  $\text{Ca}^{2+}$  binding sites on troponin C. Virtually identical reductions in force responses were also seen after exposing a fiber to 30 mM cytoplasmic  $\text{P}_i$  (142), most likely due to  $\text{Ca}^{2+}\text{-P}_i$  precipitation within the SR reducing the amount of  $\text{Ca}^{2+}$  available for rapid release (see sect. VIIA). In contrast, increasing the amount of  $\text{Ca}^{2+}$  in the SR up to approximately threefold above the endogenous level (to close to maximal capacity) did not alter the amount of  $\text{Ca}^{2+}$  released by each AP (360) (Fig. 5). This constancy in AP-induced  $\text{Ca}^{2+}$  release is evidently due to feedback effects of the released  $\text{Ca}^{2+}$  on further release (see sect. VIC8).

In summary, the SR can accumulate substantial amounts of additional  $\text{Ca}^{2+}$  without evident effect,




FIG. 5. Amount of  $\text{Ca}^{2+}$  released by single action potential (AP) stimuli versus sarcoplasmic reticulum (SR)  $\text{Ca}^{2+}$  content in a skinned fiber from rat extensor digitorum longus (EDL) muscle. The SR was progressively depleted of all its releasable  $\text{Ca}^{2+}$  in two sequences, first starting with the endogenous  $\text{Ca}^{2+}$  content and then a second time after reloading the SR to its maximal level.  $\text{Ca}^{2+}$  reuptake was blocked by an SR  $\text{Ca}^{2+}$  pump blocker during AP-induced release. The amount of  $\text{Ca}^{2+}$  released by an AP is virtually unchanged when the SR is loaded well above its endogenous content, but is decreased at lower content levels.  $\text{Ca}^{2+}$  is expressed in micromoles per liter fiber volume. [From Posterino and Lamb (360).]

whereas reduction of the amount of releasable  $\text{Ca}^{2+}$  in the SR adversely affects  $\text{Ca}^{2+}$  release and force responses.

### C. $\text{Ca}^{2+}$ Content Changes With Repeated Stimulation and With Exercise

SR  $\text{Ca}^{2+}$  content will change if there is a net influx or efflux of  $\text{Ca}^{2+}$  across the sarcolemma of a muscle fiber, and these changes in flux may affect fatigue development (64, 344, 494).  $\text{Ca}^{2+}$  may enter a muscle fiber via a number of different pathways (12), including the DHPRs (129, 260), store-operated  $\text{Ca}^{2+}$  entry (SOCE) (211, 251, 272), excitation-coupled  $\text{Ca}^{2+}$  entry (ECCE) (88, 220),  $\text{Ca}^{2+}$  leak channels (169, 216), and stretch-activated channels (SACs) (195, 448, 487). Experiments on  $^{45}\text{Ca}$  uptake in frog single fibers found a “resting”  $\text{Ca}^{2+}$  influx of  $\sim 0.25 \text{ pmol} \cdot \text{s}^{-1} \cdot \text{cm}^{-2}$  of apparent surface area, and  $\sim 1 \text{ pmol } \text{Ca}^{2+} / \text{cm}^2$  on each twitch response (in  $1.8 \text{ mM}$  extracellular  $\text{Ca}^{2+}$ ) (104). The latter equates to only  $\sim 0.7 \text{ } \mu\text{mol } \text{Ca}^{2+} / \text{kg wet wt}$  on each AP, or  $\sim 0.07\%$  of the resting SR  $\text{Ca}^{2+}$  content already present. Importantly, this is the unidirectional influx, not the net flux of  $\text{Ca}^{2+}$ . The rate of  $\text{Ca}^{2+}$  efflux in a fiber is also very low in resting fibers (time constant  $> 2 \text{ h}$ , Ref. 104), and low even with repeated activation, as is apparent from the large number of force responses that can be evoked in zero extracellular  $\text{Ca}^{2+}$  (18, 290). The combined effect of influx and efflux is that net  $\text{Ca}^{2+}$  flux is normally quite small.

Fast-twitch rat muscle stimulated in vitro at  $40 \text{ Hz}$  continuously or intermittently for a total of  $30 \text{ s}$  shows net

uptake of  $50\text{--}100 \text{ } \mu\text{mol } \text{Ca}^{2+} / \text{kg wet wt}$  (183, 313), equivalent to  $\sim 5\text{--}10\%$  of the  $\text{Ca}^{2+}$  already present in the SR endogenously and representing  $< 5\%$  of the available SR  $\text{Ca}^{2+}$  capacity. With continuous stimulation at  $1 \text{ Hz}$  for  $2\text{--}4 \text{ h}$ , fast-twitch rat muscles increase their  $\text{Ca}^{2+}$  content to  $\sim 2.5$  times the endogenous level, whereas slow-twitch muscle shows little increase (183), consistent with the endogenous  $\text{Ca}^{2+}$  and maximum SR capacity observed in such muscle types (176, 440). Frog muscle fibers stimulated in vitro to fatigue increased SR  $\text{Ca}^{2+}$  content by  $\sim 10\%$  (190). In contrast, fast- and slow-twitch muscle stimulated in situ in the rat with intermittent  $100\text{-Hz}$  tetani for  $5 \text{ min}$  showed an  $\sim 40\%$  decline in total  $\text{Ca}^{2+}$  content (286). Importantly, the latter study found no significant change in muscle  $\text{Ca}^{2+}$  content after rats had been swum to exhaustion (over  $\sim 5 \text{ min}$ ) (286). In humans running long distances, vastus lateralis muscle  $\text{Ca}^{2+}$  content increased by  $\sim 30\%$  after a  $20\text{-km}$  run, but was not significantly changed after a  $10\text{-km}$  run (338). The above findings indicate that even though muscle, particularly fast-twitch fibers, can accumulate net  $\text{Ca}^{2+}$  with repeated stimulation, there in fact was relatively little change in muscle  $\text{Ca}^{2+}$  content during exercise except when it was very prolonged.

### D. Mitochondrial $\text{Ca}^{2+}$ Handling

Mitochondria in skeletal muscle fibers can take up cytoplasmic  $\text{Ca}^{2+}$  released from the SR during twitch and tetanic responses (68, 270, 383). Such  $\text{Ca}^{2+}$  uptake may stimulate mitochondrial ATP production, although it may not be essential (68). It has been frequently suggested that  $\text{Ca}^{2+}$  uptake by the mitochondria may play a significant role in overall  $\text{Ca}^{2+}$  movements in a fiber, particularly those mitochondria situated close to the SR (223, 401), with the effect possibly being more important in frog fibers than in mammalian fibers (68, 270). However, the mitochondria do not contain high-capacity  $\text{Ca}^{2+}$  binding proteins such as calsequestrin, and the absolute amount of  $\text{Ca}^{2+}$  contained within the mitochondria in frog fibers at rest, or after stimulation by either a single tetanus or to fatigue, is  $< 2\%$  of that contained within the SR (190, 408). In humans exercising for  $1 \text{ h}$ , the amount of  $\text{Ca}^{2+}$  in the mitochondria only increased by  $\sim 50\%$  above the level in rested muscle (297). It should also be noted that the experiments in mammalian fibers reporting specific tunneling of SR  $\text{Ca}^{2+}$  to the mitochondria (223, 401) were performed with skinned fibers under conditions where the SR would have been loaded at near-maximal capacity, and this likely increased the extent of  $\text{Ca}^{2+}$  uptake by the mitochondria. In summary, the mitochondria appear to play at most a minor role in directly influencing overall  $\text{Ca}^{2+}$  movements during EC coupling.

## X. SHORTENING VELOCITY AND MECHANICAL POWER

Locomotion is generally driven by mechanical work generated by skeletal muscles, and the speed of locomotion consequently depends on the mechanical power that the muscles can produce. Mechanical power equals force times shortening velocity, and the mechanisms underlying fatigue-induced changes in isometric force production are not the same as those that affect the shortening velocity (8). This means that a decrease in maximum power output can depend mainly on decreased force production in some situations, whereas decreased shortening velocity can be of greater importance in other cases. Furthermore, decreased force production has a larger impact on the power output during movements requiring high forces, and reduced shortening velocity becomes more important as the speed of movement increases. The energy cost is considered to be higher during repeated contractions with shortening than during isometric conditions, and hence fatigue develops more rapidly when it is induced by shortening contractions (43, 102).

Fatigue-induced changes in shortening velocity can be described in terms of decreased maximal shortening velocity (i.e., velocity at zero load) and/or altered curvature of force-velocity relationship. Changes in maximal shortening velocity and curvature have been shown to develop with different time courses during fatiguing stimulation, which indicates different underlying mechanisms (228, 476).

Acidosis has classically been considered a major cause of the fatigue-induced decrease in shortening speed (8, 168). This conclusion was based on studies on frog muscle and mammalian skinned fibers studied at low temperature (10–15°C), where lowering pH to 6.2 causes a very substantial decrease in the  $F_{Ca,max}$  (>50%) and in the maximum velocity of shortening (347). However, more recent data from intact mouse type II fibers studied at 32°C show no decrease in maximal shortening velocity under acidic conditions (478). Similar results were obtained in skinned mammalian type II fibers studied at 30°C, where decreasing pH from 7.0 to 6.2 had little inhibitory effect on maximum velocity of shortening and peak power (244, 347). A larger reduction (~34%) in peak power induced by lowering pH to 6.2 at 30°C was observed in type I fibers (244), but it is very doubtful that pH ever approaches such a low level in type I fibers in normal exercise.

Increased  $P_i$  decreases the myofibrillar capacity to generate force and impairs SR  $Ca^{2+}$  handling and is therefore regarded as a major cause of fatigue (see sect. VIIA). However, the effect of increased  $P_i$  on maximum shortening velocity is very limited (99, 123, 348). Reduced ATP may result in decreased shortening speed, but ATP has to fall to very low levels (~0.5 mM) before this inhibitory

effect becomes substantial (98, 164), and this will not occur in most types of fatigue (see sect. III). Decreased  $[Ca^{2+}]_i$  results in decreased isometric force but has little effect on the maximal shortening velocity (150, 473).

Skinned fiber experiments have shown that ADP has a major inhibitory effect on maximal shortening velocity (99, 309). The free [ADP] in the myoplasm ( $[ADP]_i$ ) has been estimated to increase to no more than ~300  $\mu$ M in severely fatigued muscles, and this  $[ADP]_i$  has little impact on the maximal shortening velocity (87).  $[ADP]_i$  cannot be measured with standard biochemical methods because most ADP is bound to proteins and hence metabolically inactive (449).  $^{31}P$ -NMR spectroscopy can measure free ADP, but  $[ADP]_i$  in muscle is generally below the detection threshold and the time resolution is limited (198). Nevertheless, Hancock and co-workers (198, 199) were recently able to detect an increase in  $[ADP]_i$  to ~1.5 mM in fatigued hindlimb muscles of mice deficient of adenylate kinase (AK). AK catalyzes the reaction:  $2ADP \rightarrow ATP + AMP$ , and muscles deficient of this enzyme will therefore display larger and more long-lasting increases in  $[ADP]_i$  during fatigue than normal muscles. The unloaded shortening velocity of the gastrocnemius-plantaris-soleus muscle group of AK-deficient and wild-type mice was decreased to a similar extent during a series of repeated tetanic contraction (199). This would speak against a major role of increased  $[ADP]_i$  in the decrease in shortening velocity during fatigue, but it should be noted that the muscle group used in these experiments has a mixed fiber type composition that will complicate the interpretation of the results.

Transient increases in  $[ADP]_i$  have been suggested to occur during contractions, especially when the PCr store becomes depleted (177, 388). Accordingly, numerous studies have reported a substantial increase in IMP and a corresponding decrease in adenine nucleotides in fatigued muscles (225, 312, 390). The increase in IMP may be taken to support the occurrence of transient increases in  $[ADP]_i$  during fatiguing contractions, because the formation of IMP is driven by an increase in  $[ADP]_i$  and the flux would be very limited if  $[ADP]_i$  remained low throughout fatigue.

Experiments were designed to test the effect of the putative  $[ADP]_i$  transients on the maximal shortening velocity in single fast-twitch *Xenopus* frog and mouse muscle fibers (473, 477). The maximal shortening velocity was then first measured in a short (400 ms) tetanus. After a 2.4-s pause, it was again measured in a long (1,400 ms) tetanus, and finally after a 4-s pause in a short tetanus. The underlying assumption was that with limited PCr buffering, ADP would accumulate more in the long tetanus, and this effect would be rapidly reversed. Under control conditions, where the PCr buffering was intact and no transient increases in  $[ADP]_i$  were expected, there was no difference in the maximal shortening velocity



between the contractions. In the fatigued state, on the other hand, the maximal shortening velocity in the long tetanus was markedly lower than in the bracketing short tetani. Similar results were also obtained in unfatigued fibers where the CK reaction was inhibited by dinitrofluorobenzene (473, 477).

Results from fatigued frog and mouse muscles show a decreased curvature of the force-velocity relationship (31, 103, 476), which will limit the fatigue-induced decrease in maximum power output, whereas studies on human adductor pollicis muscles show an increased curvature (120, 228), which will further decrease the maximum power output. The mechanisms underlying these opposite results are uncertain, but they have been suggested to reflect differences in the temperature at which the experiments were performed (228). Recent studies on skinned rat fibers showed a marked decrease in the curvature of the force-velocity relationship as the temperature was increased from 15 to 30°C (123, 244). Furthermore, addition of 30 mM  $P_i$ , which has little effect on the maximum shortening velocity (see above), increased the curvature in both fast- and slow-twitch fibers when studied at 30°C but had no effect at 15°C (123).

In conclusion, fatigue involves changes in isometric force, maximal shortening velocity, and the curvature of the force-velocity relationship. These three factors have different underlying mechanisms, and they all affect the power output of fatigued muscles. Transient increases of  $[ADP]_i$  appear to have a central role in the fatigue-induced decrease in maximal shortening velocity, whereas increased  $P_i$  or  $H^+$  or decreased  $[Ca^{2+}]_i$  have little impact on this parameter.

## XI. SLOWING OF RELAXATION

Skeletal muscle fatigue is generally accompanied by a marked slowing of relaxation (Fig. 3A). This slowing may be beneficial during a prolonged isometric contraction where it increases fusion of the force output at lower stimulation frequencies, thus minimizing the force decline when the motor neuron firing rate decreases (52, 227). On the other hand, slowing of relaxation can limit performance during dynamic exercise where rapidly alternating movements are performed (8). It should also be noted that muscle cells can fatigue without any major slowing of relaxation; for instance, isolated slow-twitch fibers of mouse soleus muscles display no or very limited slowing during fatiguing stimulation (Fig. 3B) (67, 288).

Relaxation of skeletal muscle cells is a complex process that involves the following major steps: 1) SR  $Ca^{2+}$  release stops, 2)  $Ca^{2+}$  is taken up by the SR via ATP-driven pumps (myoplasmic buffers, such as parvalbumin, may contribute to the decrease in  $[Ca^{2+}]_i$  in muscles that contain such buffers and provided they are not already

saturated with  $Ca^{2+}$ ), 3) the resulting decline in  $[Ca^{2+}]_i$  means the  $Ca^{2+}$  dissociates from troponin, and 4) cross-bridge cycling ceases. Potentially, any of these steps could be slowed in fatigue and hence contribute to the slowing of relaxation. Currently available methods do not allow direct measurements of changes in the rate of each of these four steps during a fatigue experiment. In fact, it is doubtful whether any of the steps can be accurately measured during fatigue without interference from the other steps. To simplify this matter, we have developed techniques where simultaneous measurements of force and  $[Ca^{2+}]_i$  (or shortening steps during relaxation) in single muscle fibers allow us to assess the relative contribution of changes in SR  $Ca^{2+}$  handling (*steps 1 and 2*;  $Ca^{2+}$  component) and myofibrillar function (*steps 3 and 4*; cross-bridge component) (465, 478). Using these techniques, we found a slowing of SR  $Ca^{2+}$  handling in mouse FDB fibers fatigued by repeated tetani, but this was counteracted by decreased myofibrillar  $Ca^{2+}$  sensitivity, and hence, the  $Ca^{2+}$  component appeared not to affect the rate of force relaxation (465). On the other hand, when using the same experimental approaches on easily fatigued (type 1) and fatigue-resistant (type 2) *Xenopus* frog fibers, we found that both the  $Ca^{2+}$  component and the cross-bridge component contributed significantly to the fatigue-induced slowing of relaxation (478). Similar experiments cannot be performed in humans, but a recent study using human adductor pollicis muscles in vivo compared changes in the force-velocity relationship and relaxation speed during a series of fatiguing contractions (228). The results showed a strong temporal correlation between an increased curvature of the force-velocity relationship and slowed relaxation, which indicates that altered cross-bridge function contributes to the slowing of relaxation in fatigued human muscles.

In conclusion, changes in SR  $Ca^{2+}$  handling and cross-bridge function, which have the potential to cause slowed relaxation, occur in parallel during induction of fatigue. As discussed above, the relative importance of the slowing of the  $Ca^{2+}$  component and the cross-bridge component to the observed reduction in relaxation speed depends on the species and muscle studied. In addition, the relative contribution of the  $Ca^{2+}$  versus the cross-bridge component undoubtedly depends on other factors, such as the temperature and the stimulation protocol employed to induce fatigue.

Numerous studies have correlated slowing of relaxation during fatigue with concurrent metabolic changes (8, 168). A large focus has been on acidosis, but like other aspects of fatigue [i.e., isometric force (sect. VII B) and shortening velocity (sect. X)], the inhibitory effect of acidosis on the relaxation speed of mammalian muscle becomes smaller as the temperature is increased towards physiological temperatures. Nevertheless, acidification of mouse FDB fibers at ~30°C (close to the ambient tem-

perature of this superficially located muscle) caused a significant decrease in the rate of relaxation both in the unfatigued and the fatigued state (71, 478). Thus it appears that acidification contributes to the slowing of relaxation in fatigued mammalian muscle even at physiological temperatures, which agrees with *in vivo* human muscle results (77).

Changes in ATP, ADP, and  $P_i$  during fatigue will have a direct impact on both myosin and SR  $Ca^{2+}$ -ATPase, and hence, changes in these metabolites have classically been implicated in the fatigue-induced slowing of relaxation (118, 154). More recent studies show major effects of increased  $P_i$  on SR  $Ca^{2+}$  handling (sect. VIIA), and changes in ADP and ATP also affect the SR function (sect. VII C). Moreover, experiments on skinned muscle fibers, where relaxation was induced by activation of the rapid  $Ca^{2+}$  chelator diazo-2, show that both increased ADP or  $P_i$  can slow relaxation by acting on the cross bridges (217, 319). Adenylate kinase limits the accumulation of ADP during fatigue; hence, fatigued skeletal muscles of AK-deficient mice show a large increase in free [ADP] (199). Fatigued AK-deficient muscles also show more slowing of relaxation during fatigue than a wild-type muscle (199). CK-deficient muscle fibers fatigue without any significant accumulation of  $P_i$  (422). At the onset of fatigue induced by repeated tetani, wild-type muscle fibers show decreased force production and a marked slowing of relaxation, and neither of these features are seen in CK-deficient fibers (106). Thus the lack of slowing of relaxation in these support an important role of  $P_i$  in this process.

In conclusion, increases in  $H^+$ , ADP, and  $P_i$  are likely metabolic factors causing a reversible slowing of relaxation in fatigued muscles. In addition, fatigue may also induce long-lasting structural changes in SR  $Ca^{2+}$  handling and/or myofibrillar proteins that might affect the rate of relaxation (see sect. VIID).

## XII. FACTORS COUNTERACTING FATIGUE

This review mainly deals with fatigue mechanisms that cause impaired muscle function during repeated activity. However, there are also mechanisms by which repeated activity improves muscle function, and this section deals with factors that can counteract fatigue. One example of this is the fatigue-induced slowing of relaxation that will increase fusion and hence isometric force production at low stimulation frequencies. This has been shown to have an important force-preserving effect during continuous voluntary contractions, where a reduction of the  $\alpha$ -motor neuron firing frequency limits or prevents AP failure in the muscle fibers (sect. VI C). Decreased  $pH_i$  may also limit AP failure in fatigue by decreasing the  $Cl^-$  and hence the leak conductance (sect. VI C).

Muscle cells may take up water and swell during fatiguing exercise (403, 458). Muscle fiber swelling in-

creases tetanic force production and maximum shortening velocity (151) and hence counteracts the opposite changes that occur in fatigue. Interestingly, the initial beneficial effects of creatine supplementation may be due to increased force production caused by water influx (321) (see sect. VIIA5).

Submaximal force production is often increased after a brief period of muscle activity, and this phenomenon is called posttetanic potentiation (170, 366, 404). Posttetanic potentiation is generally larger in fast-twitch than in slow-twitch fibers (430). There is a correlation between this force potentiation and phosphorylation of the myosin regulatory light chain (RLC) (298, 317, 430). Furthermore, skinned fiber experiments have shown that myosin RLC phosphorylation increases force at submaximal, but not saturating,  $Ca^{2+}$  concentrations (353, 431). The increase in  $[Ca^{2+}]_i$  during contractions initiates myosin RLC phosphorylation via a  $Ca^{2+}$ /calmodulin-dependent activation of skeletal muscle myosin light-chain kinase (skMLCK) (430, 495). In addition, it appears that in fatigued muscle other factors may also have a force-potentiating effect similar to that induced by myosin RLC phosphorylation (441), possibly including oxidation effects on the contractile apparatus (255).

The increase in  $[Ca^{2+}]_i$  during contractions may also activate another  $Ca^{2+}$ -dependent kinase, calmodulin kinase II (CaMKII), which is known to regulate several proteins involved in SR  $Ca^{2+}$  handling. In fast-twitch fibers, CaMKII targets proteins involved in SR  $Ca^{2+}$  release, i.e., RyR, DHPR, and some of their associated proteins (97, 109). In slow-twitch fibers, CaMKII also acts on the SR  $Ca^{2+}$ -ATPase and phospholamban (110, 387). CaMKII has been shown to be activated in exercising humans (381). Injection of CaMKII inhibitory peptide into fast-twitch mouse FDB fibers resulted in an ~20% decrease in tetanic  $[Ca^{2+}]_i$  in unfatigued fibers, and the extent of inhibition increased with repeated contractions (436). Thus the decline in  $[Ca^{2+}]_i$  and force during fatigue might occur at a faster rate if these changes were not counteracted by  $Ca^{2+}$ -induced activation of skMLCK and CaMKII.

## XIII. RECOVERY FROM FATIGUE

In this review we consider "recovery" to start immediately after the period of fatiguing stimulation. This means that the fatigue-induced impairment in muscle function does not necessarily have to improve during the initial part of the recovery period. In fact, the opposite is frequently observed. For instance, fast-twitch frog muscle fibers show a marked force decrease after the end of fatiguing stimulation, which has been named postcontractile depression (PCD) (475) and which will be discussed below.

Depending on the type of fatigue induced, restoration of force production after fatiguing stimulation has mark-

edly different time courses. Force recovery after fatigue produced by repeated short tetani is generally completed within 30 min when tested at high frequencies, whereas the force at low-frequency stimulation may be markedly depressed for many hours. On the other hand, when fatigue is induced by continuous high-frequency stimulation, a marked force recovery occurs in the first seconds after cessation of fatiguing stimulation (sect. 1A and Fig. 2). It has been proposed that this fast component of recovery over several seconds represents the diffusion of elevated  $K^+$  from the t tubules (479). During long high-frequency tetani, reduced  $Ca^{2+}$  release was apparent in the center of a muscle fiber, consistent with  $K^+$  accumulation depolarizing the t tubules. In addition, the time course of recovery of central  $Ca^{2+}$  release and force are both consistent with the time for diffusion of small ions out of the t tubule (326) (sect. 1C).

A specific type of long-lasting lack of force recovery was originally observed by Edwards et al. (155) in human adductor pollicis muscles (155). In this type of delayed recovery from fatigue, tetanic force production is more depressed at low than at high stimulation frequencies (see sect. 1C). It should be pointed out that a greater fractional force decrease at low compared with high stimulation frequencies (low-frequency weakness) is not restricted to situations with long-lasting lack of recovery. In fact, this is seen in all situations where the lower stimulation frequency is on a steeper part of the force- $[Ca^{2+}]_i$  relationship than the higher stimulation frequency (Fig. 6). Thus all situations where tetanic  $[Ca^{2+}]_i$  and/or the myofibrillar  $Ca^{2+}$  sensitivity are reduced have the potential of showing low-frequency weakness. It can even be observed in unfatigued fibers as illustrated by experiments with inhibition of CaMKII (436). In this situation, CaMKII inhibition causes a similar reduction of tetanic  $[Ca^{2+}]_i$  at all stimulation frequencies, which results in a markedly lower force at low frequencies.

### A. Delayed Recovery From Fatigue

Following severe or prolonged exercise in humans, part of the overall force deficit can be due to changes in muscle function that persist for many hours (155, 210, 442). This deficit is not due to reduced amounts of ATP or CrP, nor to increases in associated metabolites, as these return close to resting levels within 15–60 min (22, 155, 210, 442). Furthermore, it is not due to changes in neuromuscular transmission or excitation of the sarcolemma, as the compound surface AP (M-wave) is unchanged in most circumstances (22, 155) (but see additional phenomenon of delayed tension reduction discussed below). A key feature of this long-term force reduction is that responses to low-frequency stimulation are reduced more than responses to high-frequency stimulation (e.g., 20 ver-




FIG. 6. Schematic showing mechanisms underlying a more marked force depression at low stimulation frequencies. Red arrows show what happens with a decrease in tetanic  $[Ca^{2+}]_i$ , and blue arrows with a decrease in myofibrillar  $Ca^{2+}$  sensitivity (indicated by dashed line). Note that the effect on force of both these changes is markedly larger when they originate from the steep part of the force- $[Ca^{2+}]_i$  relationship, i.e., at low stimulation frequencies.

sus 50 or 100 Hz in humans). Some studies report that the response to high-frequency stimulation is virtually unchanged (155), whereas others report that it is reduced ~20% (210, 442). The latter has been interpreted as an additional fatigue mechanism (442) and described as postcontractile depression (PCD) to link it with a phenomenon described in frog single fibers (475) (see below), although it is quite likely that the reduction in force at high frequencies is due to the same processes as that at low frequencies. For instance, a large reduction in either SR  $Ca^{2+}$  release (Fig. 7A) or the  $Ca^{2+}$  sensitivity of the contractile apparatus (Fig. 7B) could result in some force reduction even with high-frequency stimulation, as well as a proportionately larger reduction in force to low-frequency stimulation.

A long-lasting force deficit has also been observed in vitro in both fast- and slow-twitch muscles (229) and in single fast-twitch mouse fibers (89, 474). Force was reduced ~10–20% at high frequencies and >50% at low frequencies and showed little or no recovery in the period 10–60 min after stimulation. Augmenting  $Ca^{2+}$  release with caffeine fully restored the force response to high-frequency stimulation (89, 229, 474), indicating that maximum force production by the contractile apparatus was unaffected. Tetanic  $[Ca^{2+}]_i$  was measured in single mouse fibers, and the force deficit was found to be caused by a reduction in  $[Ca^{2+}]_i$  at all frequencies of stimulation,






FIG. 7. Examples of decreased force production at low-frequency stimulation observed 30 min after fatigue induced by repeated tetani in a fast-twitch mouse (A) and rat (B) FDB fiber. Note two different mechanisms: the force decrease at 50 Hz in the mouse FDB fiber can be explained by a decreased  $[Ca^{2+}]_i$  (A), and the force decrease at 30 Hz in the rat FDB fiber was not accompanied by any decrease in  $[Ca^{2+}]_i$  (B). The fact that force at 100 Hz was similar before and after fatigue shows that maximal  $Ca^{2+}$ -activated force was little affected, and hence, the force decrease at low frequencies was due to a decrease in myofibrillar  $Ca^{2+}$  sensitivity. [A from Westerblad et al. (474); B from J. D. Bruton and H. Westerblad, unpublished observations.]

whereas there were no changes in  $Ca^{2+}$  sensitivity (474). The reduction in tetanic  $[Ca^{2+}]_i$  was uniform throughout the single fiber, showing that it was not due to AP failure within the T system.

Another type of delayed force recovery, PCD, was originally observed after fatigue induced by repeated tetani in *Xenopus* fast-twitch fibers (475). PCD is manifested as a further decrease in tetanic force after cessation of fatiguing stimulation, which reaches a minimum after  $\sim 20$  min of recovery when little or no tetanic force is produced. Thereafter, force starts to recover, eventually reaching the prefatigue level, but the force recovery generally takes much longer than the recovery of ATP and

other metabolites which are completed within an hour (323). PCD is prolonged by mild mechanical stress, in particular by a brief (5 min) exposure to mildly hypotonic conditions following stimulation (69), and full recovery may then take  $\sim 16$  h (269). PCD is caused by inhibition of SR  $Ca^{2+}$  release despite normal membrane potential and AP properties (219, 266, 283, 471, 475). The exact mechanism underlying PCD has not been revealed, but it seems to involve impaired mechanical coupling between the t-tubular voltage sensors (DHPR) and the SR  $Ca^{2+}$  release channels (RyR) (69, 72).

Yet another phenomenon of long-term reduction of tetanic responses has been observed in rat (264) and human (302) muscles. In this case, tetanic force first substantially recovers after the end of stimulation and then over the next 60 min declines markedly again before subsequently recovering once more. In contrast to PCD, the force loss is seemingly attributable to a loss of muscle excitation, as is apparent from the major reduction in the M-wave that occurs in parallel (264, 302). Although failure of transmission at the neuromuscular junction has not been ruled out, this effect is probably due to failure of the sarcolemmal AP. Given the long-lasting nature of the effect, it is more likely due to changes in ionic conductances (i.e., reduced  $Na^+$  conductance or increased  $K^+$  or  $Cl^-$  conductance) rather than to changes in ion concentration gradients. The reports to date of delayed tension reduction involved experiments in which the muscle was stimulated in situ by imposed stimulation of the nerve, either continuously for  $\sim 5$  min (302) or intermittently for 15 min (264), and it is unclear whether or to what extent this effect happens after normal exercise in vivo. One study in exercising humans, however, has noted some cases of a loss in tetanic force to high-frequency stimulation that reached its maximum 1–2 h after the end of the exercise (155), which might be due to this phenomenon.

## B. Prolonged Changes in the Contractile Apparatus

Chemically skinned fibers, obtained from whole muscles fatigued in vitro by intermittent tetani, have been used to examine whether there are changes in the contractile properties of fatigued fibers that persist after the cytoplasm is replaced with a standardized solution. These experiments showed no changes in maximum force production (111, 483). In fast-twitch rodent fibers, myofibrillar  $Ca^{2+}$  sensitivity was unchanged (111), whereas slow-twitch fibers showed an overall decrease in  $Ca^{2+}$  sensitivity (111). Experiments with frog twitch fibers showed a net increase in  $Ca^{2+}$  sensitivity after fatiguing stimulation (481, 483). The cause(s) of these changes has not been identified.

### C. Prolonged Reduction in $\text{Ca}^{2+}$ Release

A number of studies have reported that  $\text{Ca}^{2+}$  release from isolated SR is reduced by ~20–40% following prolonged or intense exercise in humans (210, 284, 285) and rodents (161), although some studies found no change (111, 396) or a reduction in slow-twitch but not fast-twitch muscle (161, 221). It is unclear how long this effect persists after exercise, as this was only examined in one study to date, and the results were somewhat equivocal at the one recovery time examined (3.5 h) (210). Moreover, a major issue with such studies is that the  $\text{Ca}^{2+}$  release was triggered by nonphysiological means, such as by caffeine, chloro-*m*-cresol or the oxidizing agent  $\text{Ag}^+$ , and the rate of release was 30–1,000 times slower than occurs when activating  $\text{Ca}^{2+}$  release by the normal voltage-sensor mechanism. Given that voltage-sensor activation of  $\text{Ca}^{2+}$  release is frequently found to be unaffected by conditions that inhibit  $\text{Ca}^{2+}$  release in isolated SR channels (e.g., raised  $[\text{H}^+]$ , lactate, oxidation), the above findings cannot be taken as strong evidence that exercise-induced changes in the release channels themselves are the primary cause of either short-term or long-term reductions in  $\text{Ca}^{2+}$  release following exercise.

The long-term reduction in AP-induced  $\text{Ca}^{2+}$  release in isolated murine fibers (474) becomes more pronounced if the rise in  $[\text{Ca}^{2+}]_i$  during the fatiguing stimulation is prolonged or pharmacologically augmented, and repeated fatiguing bouts have a cumulative effect (89, 92). Potentiating tetanic  $[\text{Ca}^{2+}]_i$  also greatly prolongs PCD in *Xenopus* fast-twitch fibers (69, 70, 269). On the other hand, in a recent study fatigue was induced by repeated tetani in isolated mouse fibers exposed to *N*-benzyl-*p*-toluene sulfonamide (BTS), which inhibits cross-bridge force production (66). The long-term depression of SR  $\text{Ca}^{2+}$  release was no more marked in BTS-exposed fibers compared with control fibers despite the fact that the former experienced a markedly larger  $[\text{Ca}^{2+}]_i$ -time integral during induction of fatigue.

Experiments in skinned fibers with functional EC coupling show that coupling is disrupted in both mammalian and amphibian fibers if the cytoplasmic  $[\text{Ca}^{2+}]$  is raised to tetanic levels (2–10  $\mu\text{M}$ ) for a prolonged period or to higher levels even briefly (254, 450). The disruption occurs irrespective of whether the  $\text{Ca}^{2+}$  is applied exogenously or released from the SR by triggering the normal release mechanism, and it can also be induced by triggering excessive  $\text{Ca}^{2+}$  release whilst the fibers are intact (254, 450). The effect is caused primarily by relatively high  $[\text{Ca}^{2+}]$  near the release channels rather than by prolonged small rises in resting  $[\text{Ca}^{2+}]$  (450). The reduced responses are not due to depolarization of the T system, or to dysfunction of the release channels, which can still be potently activated by direct stimulation. Instead, the effect appears to be due to the voltage sensors failing to

properly activate the release channels, quite possibly because the triad junctions become distorted following the  $\text{Ca}^{2+}$  exposure (254). The elevated  $[\text{Ca}^{2+}]$  appears to induce its effect via a pH-dependent enzymatic reaction that does not involve phosphorylation or oxidation (254), but the exact mechanism is uncertain.

It has frequently been suggested that  $\text{Ca}^{2+}$ -dependent proteolysis may play a role in muscle fatigue and damage (40, 41, 76, 182). Skeletal muscle contains the ubiquitous calcium-dependent neutral proteases  $\mu$ -calpain and m-calpain, as well as a muscle-specific isoform, calpain-3 (186). Much of the ubiquitous calpain is freely diffusible, but some is normally bound at triad junction (182), and raising  $[\text{Ca}^{2+}]$  within the physiological range increases calpain binding there (322, 367).  $\text{Ca}^{2+}$ -dependent uncoupling is prevented in toad skinned fiber by leupeptin, a calpain inhibitor, but only at low  $[\text{Ca}^{2+}]$  when the uncoupling proceeds slowly (254, 451). Calpain inhibitors, however, were not found to prevent either long-duration fatigue in mouse fibers (89) or hypotonic-induced PCD in frog fibers (69). Calpain inhibitors such as leupeptin are quite poor at inhibiting in situ calpain-dependent proteolysis of structural proteins (141), unless used at very high concentration (241, 322, 451), so it still seems possible that they might play a role in the  $\text{Ca}^{2+}$ -dependent uncoupling. It is known, however, that  $\text{Ca}^{2+}$ -dependent uncoupling does not involve proteolysis of the DHPRs, RyRs, or another junctional protein, triadin (254). Interestingly, triad junction structure has been found to depend both on calpain-susceptible and calpain-resistant interactions (242), so  $\text{Ca}^{2+}$ -dependent disruption of triad structure and coupling may involve a number of different processes, leading to quite complex behavior of onset and recovery. Finally, neither  $\mu$ -calpain nor calpain-3 was detectably activated by exhaustive sprint cycling in untrained human subjects or by prolonged cycling in trained subjects (320). Possibly calpain activation may only occur with relatively extreme exercise or when there is a damaging eccentric component.

In summary, the delayed recovery after fatigue is due primarily to a reduction in SR  $\text{Ca}^{2+}$  release, which may involve elevated  $[\text{Ca}^{2+}]_i$  in some way causing impaired coupling between the t-tubular voltage sensors and the SR  $\text{Ca}^{2+}$  release channels, but the exact mechanisms involved remain uncertain.

### XIV. BLOOD FLOW AND INTRACELLULAR $\text{Po}_2$

In intact muscles, the supply of  $\text{O}_2$  depends on blood flow and the diffusion of  $\text{O}_2$  from capillary blood across the interstitial space and into the muscle fiber. A long-debated issue is whether  $\text{O}_2$  supply through this pathway is adequate or whether there are conditions in which  $\text{O}_2$  supply is limited and contributes to reduced muscle per-

formance and more rapid fatigue (429). It has already been noted that blood flow ceases in a continuous maximal contraction (33), and under these circumstances, failure of  $O_2$  supply will presumably contribute to the rapid fatigue. The dramatic effect of preventing oxidative phosphorylation with cyanide is illustrated in Figure 1.

During intermittent contraction, blood flow recovers between contractions, and it might be assumed that autoregulation of the blood supply would match the blood flow to the overall metabolic needs of the muscle. However, there is evidence that this does not necessarily occur. For instance, simply raising the arm above the head accelerates the rate of fatigue of the adductor pollicis (490). This arises because the hydrostatic fall in arterial pressure in the hand is not matched by the fall in venous pressure presumably because the veins collapse. Thus a small fall in driving pressure occurs in the raised hand, which presumably reduces the blood flow.

Some athletes show a mild arterial hypoxemia during high-intensity exercise, and this is associated with a reduced performance (380). This performance decline is partly caused by peripheral fatigue and can be reduced by an increased inspired  $[O_2]$  and exacerbated by reduced inspired  $[O_2]$  (13, 215). Furthermore, blood doping and abuse of recombinant human erythropoietin have become an increasing problem in endurance sports because they enhance maximum oxygen uptake and improve aerobic performance (126, 156).

The development of a NMR method for measuring oxygenated versus deoxygenated myoglobin in muscle has allowed estimates of the muscle  $P_{O_2}$  and provided new insights into the diffusion of  $O_2$  from capillaries to the muscle fibers. The  $P_{O_2}$  in resting capillaries and muscle are respectively 44 and 34 mmHg, whereas during exercise these values fall to 34 and 3 mmHg (376). The average value of 3 mmHg does not change in muscle from mild to severe exercise, suggesting that the diffusion gradient is maximal and that further increases in  $O_2$  supply must come from increasing blood flow, opening new capillaries, etc. (377).

In summary, in intact individuals, supply of  $O_2$  from capillaries to muscle fibers provides one limit to muscle performance during prolonged aerobic exercise.

## XV. CONCLUDING REMARKS

It is widely accepted that fatigue has many causes and that in different physical activities both the mechanisms involved and their quantitative importance can vary. Consider the fatigue caused by voluntary repeated short tetani, which in a human can lead to a 50% reduction in electrically stimulated force (48). Of the 50% decline, we suggest that the early fall (phase 1 in Fig. 1), which is usually a  $\sim 10\%$  reduction, is likely to be caused by a

reduction in  $F_{Ca,max}$  due to increased  $P_i$ . The remaining 40% has to be attributed to changes in activation, which include contributions from decreased  $Ca^{2+}$  sensitivity of the contractile proteins and reduced SR  $Ca^{2+}$  release. The reduced  $Ca^{2+}$  sensitivity could have contributions both from metabolites such as  $P_i$  and from the effects of ROS. The most likely causes for reduced SR  $Ca^{2+}$  release appear to be precipitation of  $Ca^{2+}$  phosphate in the SR with a contribution from reduced ATP and raised  $Mg^{2+}$ . Our interpretation of the evidence is that failure of the AP in the surface or the t-tubular membrane probably makes little contribution to the failure of SR  $Ca^{2+}$  release during repeated short tetani. If failure of the AP does contribute to fatigue, it is most likely to be during continuous maximal contractions when absence of perfusion enhances the  $K^+$  accumulation, although slowing of the cortical firing rate, acid-induced reduction of the  $Cl^-$  conductance, acceleration of the  $Na^+-K^+$  pump, and the large safety factor of the AP all help to minimize such failure. Figure 8 summarizes the mechanisms we believe to be important in fatigue.

The evidence for the above mechanisms comes largely from experiments on isolated animal muscle preparations. In intact muscle, the force reduction is easy to measure, and many of the metabolic changes can be estimated from NMR or muscle biopsies, but this information does not allow identification of the mechanisms involved. One promising new approach to study fatigue in intact animals is the use of genetically encoded calcium sensors that can provide  $Ca^{2+}$  signals from intact muscles (382, 434). This method could be used to assess the importance of failure of  $Ca^{2+}$  release in an intact, perfused model of fatigue. Another possibility would be if the target protein for the ROS-induced reduction in  $Ca^{2+}$  sensitivity was identified; this might allow assessment of this form of fatigue from a muscle biopsy. Other molecular targets, e.g., phosphorylation of the RyR (460), are also susceptible to this approach.

A major focus for the future must be to identify the mechanisms that contribute to human fatigue in various activities and particularly during disease processes. Increased fatigability is reported in numerous clinical conditions; in fact, it has been claimed to be the most frequent presenting symptom in a primary healthcare setting (205). Fatigue, as reported by patients, would occur when the intended physical activity can no longer be continued or is perceived as involving excessive effort and discomfort. Thus the decreased fatigue resistance in many clinical conditions can be due to other factors than a reduced fatigue resistance of the muscle cells per se, and this is illustrated in Figure 9. In principle, the time until the intended force or power output no longer can be maintained depends on the interaction between the required force, the maximum force that the muscles can produce, and the endurance of the muscle cells. For instance,






FIG. 8. Schematic diagram illustrating the major mechanisms that contribute to muscle fatigue. Heading in each box identifies subcellular function, and the subsequent list indicates cellular changes occurring during fatigue that influence the subcellular function. SM, surface membrane; TT, t tubule; SR, sarcoplasmic reticulum; AP, action potential.

obesity is associated with increased fatigability during activities where counteracting gravity has a central role (e.g., walking and running), and this can be explained by an increase in the required force due to the increased body weight. On the other hand, obesity has little effect on the performance during activities where gravity is less important (e.g., swimming).

Numerous disorders like cancer cachexia, general inflammatory diseases, sepsis, burns, human immunodeficiency syndrome, chronic kidney failure, muscular dystrophies, as well as normal aging are associated with a loss of skeletal muscle tissue (291). The decreased muscle mass in these conditions results in a decreased capacity of muscles to generate force. Early fatigue development can then occur because muscles always have to work at a higher fraction of their maximal capacity, whereas the actual fatigue resistance of the muscle cells may not be affected.

There are also disorders where the fatigue resistance of the muscle cells is decreased, but this group appears rather small. Mutations in proteins involved in energy metabolism can result in a markedly increased fatigability; for instance, patients with myophosphorylase deficiency cannot break down glycogen and show a markedly accelerated fatigue development (78). On the other hand, myoadenylate deaminase deficiency, which decreases the ability to deaminate AMP to IMP, has a relatively high incidence (~2%) in the general population and is only associated with limited changes in muscle performance (122, 336).

Patients with mitochondrial myopathies and deficient respiratory chain function may display severe muscle dysfunction (128). Intuitively, mitochondrial myopathy would lead to premature fatigue development due to an increased dependency on anaerobic metabolism during fatiguing stimulation. However, a recent study em-




FIG. 9. Schematic to illustrate different mechanisms leading to exhaustion. Dashed line shows how the maximum force (or power) declines during repeated tetani. Solid red line indicates a submaximal force required for a particular activity. Exhaustion (failure to produce the required force) occurs at the intersection of the two lines. Increases and decreases in the required force (arrow 1) will cause earlier and later onset of exhaustion, respectively. Increases and decreases in the maximum force that the muscle can produce (arrow 2) will also change the time to exhaustion. Finally, changes in the intrinsic fatigability of the muscle (arrow 3) will also change the time to exhaustion.

employing a mouse model for mitochondrial myopathy, induced by skeletal muscle specific disruption of the gene for mitochondrial transcription factor A, showed no increase in the rate of fatigue development compared with wild-type controls, but the ability to generate force was markedly decreased in mitochondrial myopathy muscles (489). The unexpected lack of effect on fatigue development could be explained by an increased mitochondrial mass in mitochondrial myopathy muscle that compensated for the respiratory chain deficiency. Thus reduced mitochondrial ATP production might not be as critical for the pathophysiology of mitochondrial myopathy as previously thought.

Patients with congestive heart failure (CHF) frequently report decreased fatigue resistance, which intuitively would be related to an impaired  $O_2$  delivery to the working muscles. However, there is a poor correlation between the decrease in fatigue resistance and the decrease in heart function (486). This suggests that CHF causes intrinsic defects in skeletal muscles, and numerous studies have reported important functional and biochemical changes in skeletal muscle cells associated with CHF (for recent reviews, see Refs. 202, 289, 330). These changes may affect fatigue development by acting both on myofibrillar function and on SR  $Ca^{2+}$  handling (287, 288, 372, 457).

In the clinical setting, a careful analysis of the muscle function is important for designing optimal strategies to treat patients with decreased fatigue resistance. As described above, increased fatigability is frequently due to

a general decrease in muscle force production rather than a decreased fatigue resistance of the muscle cells. This means that a strength training program to increase, or avoid further loss of, muscle mass may improve fatigue resistance more effectively than an endurance training program.

#### ACKNOWLEDGMENTS

Address for reprint requests and other correspondence: D. G. Allen, School of Medical Sciences and Bosch Institute, Univ. of Sydney F13, NSW 2006, Australia (e-mail: davida@physiol.usyd.edu.au).

#### GRANTS

We thank the Australian National Health and Medical Research Council, the Australian Research Council, the Swedish Research Council, the Karolinska Institute, and the Swedish National Center for Sports Research for research funding.

#### REFERENCES

- Abbate F, Bruton JD, De Haan A, Westerblad H. Prolonged force increase following a high-frequency burst is not due to a sustained elevation of  $[Ca^{2+}]_i$ . *Am J Physiol Cell Physiol* 283: C42–C47, 2002.
- Adams GR, Fisher MJ, Meyer RA. Hypercapnic acidosis and increased  $H_2PO_4^-$  concentrations do not decrease force in cat skeletal muscle. *Am J Physiol Cell Physiol* 260: C805–C812, 1991.
- Adrian RH, Bryant SH. On the repetitive discharge in myotonic muscle fibres. *J Physiol* 240: 505–515, 1974.
- Adrian RH, Chandler WK, Rakowski RF. Charge movement and mechanical repriming in skeletal muscle. *J Physiol* 254: 361–388, 1976.
- Adrian RH, Peachey LD. Reconstruction of the action potential of frog sartorius muscle. *J Physiol* 235: 103–131, 1973.
- Ahern GP, Laver DR. ATP inhibition and rectification of a  $Ca^{2+}$ -activated anion channel in sarcoplasmic reticulum of skeletal muscle. *Biophys J* 74: 2335–2351, 1998.
- Allen DG, Duty S, Westerblad H. The activation of central myofibrils during muscle fatigue caused by repeated short tetani. *J Muscle Res Cell Motil* 14: 543–545, 1993.
- Allen DG, Lännergren J, Westerblad H. Muscle cell function during prolonged activity: cellular mechanisms of fatigue. *Exp Physiol* 80: 497–527, 1995.
- Allen DG, Lee JA, Westerblad H. Intracellular calcium and tension during fatigue in isolated single muscle fibers from *Xenopus laevis*. *J Physiol* 415: 433–458, 1989.
- Allen DG, Orchard CH. Myocardial contractile function during ischemia and hypoxia. *Circ Res* 60: 153–168, 1987.
- Allen DG, Westerblad H. Role of phosphate and calcium stores in muscle fatigue. *J Physiol* 536: 657–665, 2001.
- Allen DG, Whitehead NP, Yeung EW. Mechanisms of stretch-induced muscle damage in normal and dystrophic muscle: role of ionic changes. *J Physiol* 567: 723–735, 2005.
- Amann M, Romer LM, Pegelow DF, Jacques AJ, Hess CJ, Dempsey JA. Effects of arterial oxygen content on peripheral locomotor muscle fatigue. *J Appl Physiol* 101: 119–127, 2006.
- Andrade FH, Reid MB, Allen DG, Westerblad H. Effect of hydrogen peroxide and dithiothreitol on contractile function of single skeletal muscle fibres from the mouse. *J Physiol* 509: 565–575, 1998.
- Andrade FH, Reid MB, Westerblad H. Contractile response of skeletal muscle to low peroxide concentrations: myofibrillar calcium sensitivity as a likely target for redox-modulation. *FASEB J* 15: 309–311, 2001.
- Andrews MA, Godt RE, Nosek TM. Influence of physiological  $L(+)$ -lactate concentrations on contractility of skinned striated muscle fibers of rabbit. *J Appl Physiol* 80: 2060–2065, 1996.

17. **Arbogast S, Reid MB.** Oxidant activity in skeletal muscle fibers is influenced by temperature, CO<sub>2</sub> level and muscle-derived nitric oxide. *Am J Physiol Regul Integr Comp Physiol* 287: R698–R705, 2004.
18. **Armstrong CM, Bezanilla FM, Horowicz P.** Twitches in the presence of ethylene glycol bis-(aminoethyl ether)-N,N'-tetracetic acid. *Biochim Biophys Acta* 267: 605–608, 1972.
19. **Ashley CC, Mulligan IP, Lea TJ.** Ca<sup>2+</sup> and activation mechanisms in skeletal muscle. *Q Rev Biophys* 24: 1–73, 1991.
20. **Baker AJ, Brandes R, Weiner MW.** Effects of intracellular acidosis on Ca<sup>2+</sup> activation, contraction, and relaxation of frog skeletal muscle. *Am J Physiol Cell Physiol* 268: C55–C63, 1995.
21. **Baker AJ, Carson PJ, Miller RG, Weiner MW.** Metabolic and nonmetabolic components of fatigue monitored with <sup>31</sup>P-NMR. *Muscle Nerve* 17: 1002–1009, 1994.
22. **Baker AJ, Kostov KG, Miller RG, Weiner MW.** Slow force recovery after long-duration exercise: metabolic and activation factors in muscle fatigue. *J Appl Physiol* 74: 2294–2300, 1993.
23. **Baker AJ, Languemare MC, Brandes R, Weiner MW.** Intracellular tetanic calcium signals are reduced in fatigue of whole skeletal muscle. *Am J Physiol Cell Physiol* 264: C577–C582, 1993.
24. **Baldwin KM, Klinkerfuss GH, Terjung RL, Mole PA, Holloszy JO.** Respiratory capacity of white, red, and intermediate muscle: adaptive response to exercise. *Am J Physiol* 222: 373–378, 1972.
25. **Balog EM, Fitts RH.** Effects of fatiguing stimulation on intracellular Na<sup>+</sup> and K<sup>+</sup> in frog skeletal muscle. *J Appl Physiol* 81: 679–685, 1996.
26. **Balog EM, Fitts RH.** Effects of depolarization and low intracellular pH on charge movement currents of frog skeletal muscle fibers. *J Appl Physiol* 90: 228–234, 2001.
27. **Balog EM, Fruen BR, Kane PK, Louis CF.** Mechanisms of P<sub>i</sub> regulation of the skeletal muscle SR Ca<sup>2+</sup> release channel. *Am J Physiol Cell Physiol* 278: C601–C611, 2000.
28. **Balog EM, Thompson LV, Fitts RH.** Role of sarcolemma action potentials and excitability in muscle fatigue. *J Appl Physiol* 76: 2157–2162, 1994.
29. **Bangsbo J, Graham T, Johansen L, Strange S, Christensen C, Saltin B.** Elevated muscle acidity and energy production during exhaustive exercise in humans. *Am J Physiol Regul Integr Comp Physiol* 263: R891–R899, 1992.
30. **Bangsbo J, Madsen K, Kiens B, Richter EA.** Effect of muscle acidity on muscle metabolism and fatigue during intense exercise in man. *J Physiol* 495: 587–596, 1996.
31. **Barclay CJ.** Mechanical efficiency and fatigue of fast and slow muscles of the mouse. *J Physiol* 497: 781–794, 1996.
32. **Barclay CJ.** Modelling diffusive O<sub>2</sub> supply to isolated preparations of mammalian skeletal and cardiac muscle. *J Muscle Res Cell Motil* 26: 225–235, 2005.
33. **Barcroft H, Millen JL.** The blood flow through muscle during sustained contraction. *J Physiol* 97: 17–31, 1939.
34. **Barnes M, Gibson LM, Stephenson DG.** Increased muscle glycogen content is associated with increased capacity to respond to T-system depolarisation in mechanically skinned skeletal muscle fibres from the rat. *Pflügers Arch* 442: 101–106, 2001.
35. **Baylor SM, Chandler WK, Marshall MW.** Sarcoplasmic reticulum calcium release in frog skeletal muscle fibres estimated from Arsenazo III calcium transients. *J Physiol* 344: 625–666, 1983.
36. **Baylor SM, Hollingworth S.** Model of sarcomeric Ca<sup>2+</sup> movements, including ATP Ca<sup>2+</sup> binding and diffusion, during activation of frog skeletal muscle. *J Gen Physiol* 112: 297–316, 1998.
37. **Baylor SM, Hollingworth S.** Sarcoplasmic reticulum calcium release compared in slow-twitch and fast-twitch fibres of mouse muscle. *J Physiol* 551: 125–138, 2003.
38. **Beard NA, Laver DR, Dulhunty AF.** Calsequestrin and the calcium release channel of skeletal and cardiac muscle. *Prog Biophys Mol Biol* 85: 33–69, 2004.
39. **Beckman JS, Beckman TW, Chen J, Marshall PA, Freeman BA.** Apparent hydroxyl radical production by peroxynitrite: implications for endothelial injury from nitric oxide and superoxide. *Proc Natl Acad Sci USA* 87: 1620–1624, 1990.
40. **Belcastro AN.** Skeletal muscle calcium-activated neutral protease (calpain) with exercise. *J Appl Physiol* 74: 1381–1386, 1993.
41. **Belcastro AN, Shewchuk LD, Raj DA.** Exercise-induced muscle injury: a calpain hypothesis. *Mol Cell Biochem* 179: 135–145, 1998.
42. **Bellemare F, Woods JJ, Johansson R, Bigland-Ritchie B.** Motor-unit discharge rates in maximal voluntary contractions of three human muscles. *J Neurophysiol* 50: 1380–1392, 1983.
43. **Beltman JG, Van Der Vliet MR, Sargeant AJ, De Haan A.** Metabolic cost of lengthening, isometric and shortening contractions in maximally stimulated rat skeletal muscle. *Acta Physiol Scand* 182: 179–187, 2004.
44. **Bemben MG, Lamont HS.** Creatine supplementation and exercise performance: recent findings. *Sports Med* 35: 107–125, 2005.
45. **Berchtold MW, Brinkmeier H, Muntenner M.** Calcium ion in skeletal muscle: its crucial role for muscle function, plasticity, and disease. *Physiol Rev* 80: 1215–1265, 2000.
46. **Bergström J, Hermansen L, Hultman E, Saltin B.** Diet, muscle glycogen and physical performance. *Acta Physiol Scand* 71: 140–150, 1967.
47. **Bezanilla F, Caputo C, Gonzalez-Serratos H, Venosa RA.** Sodium dependence of the inward spread of activation in isolated twitch muscles of the frog. *J Physiol* 223: 507–523, 1972.
48. **Bigland-Ritchie B, Cafarelli E, Vøllestad NK.** Fatigue of submaximal static contractions. *Acta Physiol Scand Suppl* 556: 137–148, 1986.
49. **Bigland-Ritchie B, Furbush F, Woods JJ.** Fatigue of intermittent submaximal voluntary contractions: central and peripheral factors. *J Appl Physiol* 61: 421–429, 1986.
50. **Bigland-Ritchie B, Johansson R, Lippold OC, Smith S, Woods JJ.** Changes in motoneuron firing rates during sustained maximal voluntary contractions. *J Physiol* 340: 335–346, 1983.
51. **Bigland-Ritchie B, Johansson R, Lippold OC, Woods JJ.** Contractile speed and EMG changes during fatigue of sustained maximal voluntary contractions. *J Neurophysiol* 50: 313–324, 1983.
52. **Bigland-Ritchie B, Jones DA, Woods JJ.** Excitation frequency and muscle fatigue: electrical responses during human voluntary and stimulated contractions. *Exp Neurol* 64: 414–427, 1979.
53. **Bigland-Ritchie B, Woods JJ.** Changes in muscle contractile properties and neural control during human muscular fatigue. *Muscle Nerve* 7: 691–699, 1984.
54. **Bigland-Ritchie B, Zijdevind I, Thomas CK.** Muscle fatigue induced by stimulation with and without doublets. *Muscle Nerve* 23: 1348–1355, 2000.
55. **Blazev R, Lamb GD.** Adenosine inhibits depolarization-induced Ca<sup>2+</sup> release in mammalian skeletal muscle. *Muscle Nerve* 22: 1674–1683, 1999.
56. **Blazev R, Lamb GD.** Low [ATP] and elevated [Mg<sup>2+</sup>] reduce depolarization-induced Ca<sup>2+</sup> release in mammalian skeletal muscle. *J Physiol* 520: 203–215, 1999.
57. **Blijham PJ, Hengstman GJ, Ter Laak HJ, Van Engelen BG, Zwarts MJ.** Muscle-fiber conduction velocity and electromyography as diagnostic tools in patients with suspected inflammatory myopathy: a prospective study. *Muscle Nerve* 29: 46–50, 2004.
58. **Booth J, Wilmshire BR, MacDonald AD, Zeyl A, McGhee S, Calvert D, Marino FE, Storlien LH, Taylor NA.** Whole-body pre-cooling does not alter human muscle metabolism during submaximal exercise in the heat. *Eur J Appl Physiol* 84: 587–590, 2001.
59. **Bortolotto SK, Cellini M, Stephenson DG, Stephenson GM.** MHC isoform composition and Ca<sup>2+</sup>- or Sr<sup>2+</sup>-activation properties of rat skeletal muscle fibers. *Am J Physiol Cell Physiol* 279: C1564–C1577, 2000.
60. **Bottinelli R.** Functional heterogeneity of mammalian single muscle fibres: do myosin isoforms tell the whole story? *Pflügers Arch* 443: 6–17, 2001.
61. **Bottinelli R, Reggiani C.** Human skeletal muscle fibres: molecular and functional diversity. *Prog Biophys Mol Biol* 73: 195–262, 2000.
62. **Bretag AH.** Muscle chloride channels. *Physiol Rev* 67: 618–724, 1987.
63. **Brotto MA, Van Leyen AS, Nosek CM, Brotto LS, Nosek TM.** Hypoxia and fatigue-induced modification of function and proteins in intact and skinned murine diaphragm muscle. *Pflügers Arch* 440: 727–734, 2000.
64. **Brotto MA, Nagaraj RY, Brotto LS, Takeshima H, Ma JJ, Nosek TM.** Defective maintenance of intracellular Ca<sup>2+</sup> homeo-



- stasis is linked to increased muscle fatigability in the MG29 null mice. *Cell Res* 14: 373–378, 2004.
65. **Brotto MA, Nosek TM.** Hydrogen peroxide disrupts  $\text{Ca}^{2+}$  release from the sarcoplasmic reticulum of rat skeletal muscle fibers. *J Appl Physiol* 81: 731–737, 1996.
  66. **Bruton J, Pinniger GJ, Lännergren J, Westerblad H.** The effects of the myosin-II inhibitor *N*-benzyl-*p*-toluene sulphonamide on fatigue in mouse single intact toe muscle fibres. *Acta Physiol* 186: 59–66, 2006.
  67. **Bruton J, Tavi P, Aydin J, Westerblad H, Lännergren J.** Mitochondrial and myoplasmic  $[\text{Ca}^{2+}]$  in single fibres from mouse limb muscles during repeated tetanic contractions. *J Physiol* 551: 179–190, 2003.
  68. **Bruton JD, Dahlstedt AJ, Abbate F, Westerblad H.** Mitochondrial function in intact skeletal muscle fibres of creatine kinase deficient mice. *J Physiol* 552: 393–402, 2003.
  69. **Bruton JD, Lännergren J, Westerblad H.** Mechano-sensitive linkage in excitation-contraction coupling in frog skeletal muscle. *J Physiol* 484: 737–742, 1995.
  70. **Bruton JD, Lännergren J, Westerblad H.** Effects of repetitive tetanic stimulation at long intervals on excitation-contraction coupling in frog skeletal muscle. *J Physiol* 495: 15–22, 1996.
  71. **Bruton JD, Lännergren J, Westerblad H.** Effects of  $\text{CO}_2$ -induced acidification on the fatigue resistance of single mouse muscle fibers at 28 degrees C. *J Appl Physiol* 85: 478–483, 1998.
  72. **Bruton JD, Szentesi P, Lännergren J, Westerblad H, Kovacs L, Csernoch L.** Frog skeletal muscle fibers recovering from fatigue have reduced charge movement. *J Muscle Res Cell Motil* 21: 621–628, 2000.
  73. **Bruton JD, Westerblad H, Katz A, Lännergren J.** Augmented force output in skeletal muscle fibres of *Xenopus* following a preceding bout of activity. *J Physiol* 493: 211–217, 1996.
  74. **Bruton JD, Wretman C, Katz A, Westerblad H.** Increased tetanic force and reduced myoplasmic  $[\text{P}_i]$  following a brief series of tetani in mouse soleus muscle. *Am J Physiol Cell Physiol* 272: C870–C874, 1997.
  75. **Burke RE, Levine DN, Trairris P, Zajac FE.** Physiological types and histochemical profiles in muscle motor units of the cat gastrocnemius. *J Physiol* 234: 723–748, 1973.
  76. **Byrd SK.** Alterations in the sarcoplasmic reticulum: a possible link to exercise-induced muscle damage. *Med Sci Sports Exercise* 24: 531–536, 1992.
  77. **Cady EB, Elshove H, Jones DA, Moll A.** The metabolic causes of slow relaxation in fatigued human skeletal muscle. *Working Database* 418: 327–337, 1989.
  78. **Cady EB, Jones DA, Lynn J, Newham DJ.** Changes in force and intracellular metabolites during fatigue of human skeletal muscle. *J Physiol* 418: 311–325, 1989.
  79. **Cairns SP.** Lactic acid and exercise performance: culprit or friend? *Sports Med* 36: 279–291, 2006.
  80. **Cairns SP, Buller SJ, Loiselle DS, Renaud JM.** Changes of action potentials and force at lowered  $[\text{Na}]_o$  in mouse skeletal muscle: implications for fatigue. *Am J Physiol Cell Physiol* 285: C1131–C1141, 2003.
  81. **Cairns SP, Dulhunty AF.** High-frequency fatigue in rat skeletal muscle: role of extracellular ion concentrations. *Muscle Nerve* 18: 890–898, 1995.
  82. **Cairns SP, Hing WA, Slack JR, Mills RG, Loiselle DS.** Different effects of raised  $[\text{K}^+]$  on membrane potential and contraction in fast- and slow-twitch muscle. *Am J Physiol Cell Physiol* 273: C598–C611, 1997.
  83. **Cairns SP, Ruzhynsky V, Renaud JM.** Protective role of extracellular chloride in fatigue of isolated mammalian skeletal muscle. *Am J Physiol Cell Physiol* 287: C762–C770, 2004.
  84. **Callahan LA, She ZW, Nosek TM.** Superoxide, hydroxyl radical, hydrogen peroxide effects on single-diaphragm fiber contractile apparatus. *J Appl Physiol* 90: 45–54, 2001.
  85. **Carlson FD.** *Muscle Physiology*, edited by Wilkie DR. Englewood Cliffs, NJ: Prentice Hall, 1974.
  86. **Chase PB, Kushmerick MJ.** Effects of pH on contraction of rabbit fast and slow skeletal muscle fibers. *Biophys J* 53: 935–946, 1988.
  87. **Chase PB, Kushmerick MJ.** Effect of physiological ADP concentrations on contraction of single skinned fibers from rabbit fast and slow muscles. *Am J Physiol Cell Physiol* 268: C480–C489, 1995.
  88. **Cherednichenko G, Hurne AM, Fessenden JD, Lee EH, Allen PD, Beam KG, Pessah IN.** Conformational activation of  $\text{Ca}^{2+}$  entry by depolarization of skeletal myotubes. *Proc Natl Acad Sci USA* 101: 15793–15798, 2004.
  89. **Chin ER, Allen DG.** The role of elevations in intracellular  $\text{Ca}^{2+}$  concentration in the development of low frequency fatigue in mouse single muscle fibres. *J Physiol* 491: 813–824, 1996.
  90. **Chin ER, Allen DG.** Effects of reduced muscle glycogen concentration on force,  $\text{Ca}^{2+}$  release and contractile protein function in intact mouse skeletal muscle. *J Physiol* 498: 17–29, 1997.
  91. **Chin ER, Allen DG.** The contribution of pH-dependent mechanisms to fatigue at different intensities in mammalian single muscle fibres. *J Physiol* 512: 831–840, 1998.
  92. **Chin ER, Balnave CD, Allen DG.** Role of intracellular calcium and metabolites in low-frequency fatigue in mouse skeletal muscle. *Am J Physiol Cell Physiol* 272: C550–C559, 1997.
  93. **Chua M, Dulhunty AF.** Inactivation of excitation-contraction coupling in rat extensor digitorum longus and soleus muscles. *J Gen Physiol* 91: 737–757, 1988.
  94. **Clanton TL, Zuo L, Klawitter P.** Oxidants and skeletal muscle function: physiologic and pathophysiologic implications. *Proc Soc Exp Biol Med* 222: 253–262, 1999.
  95. **Clausen T.**  $\text{Na}^+/\text{K}^+$  pump regulation and skeletal muscle contractility. *Physiol Rev* 83: 1269–1324, 2003.
  96. **Clausen T, Overgaard K, Nielsen OB.** Evidence that the  $\text{Na}^+/\text{K}^+$  leak/pump ratio contributes to the difference in endurance between fast- and slow-twitch muscles. *Acta Physiol Scand* 180: 209–216, 2004.
  97. **Colpo P, Nori A, Sacchetto R, Damiani E, Margreth A.** Phosphorylation of the triadin cytoplasmic domain by CaM protein kinase in rabbit fast-twitch muscle sarcoplasmic reticulum. *Mol Cell Biochem* 223: 139–145, 2001.
  98. **Cooke R, Bialek W.** Contraction of glycerinated muscle fibers as a function of the ATP concentration. *Biophys J* 28: 241–258, 1979.
  99. **Cooke R, Pate E.** The effects of ADP and phosphate on the contraction of muscle fibers. *Biophys J* 48: 789–798, 1985.
  100. **Coonan JR, Lamb GD.** Effect of transverse-tubular chloride conductance on excitability in skinned skeletal muscle fibres of rat and toad. *J Physiol* 509: 551–564, 1998.
  101. **Coupland ME, Puchert E, Ranatunga KW.** Temperature dependence of active tension in mammalian (rabbit psoas) muscle fibres: effect of inorganic phosphate. *J Physiol* 536: 879–891, 2001.
  102. **Cummins ME, Soomal RS, Curtin NA.** Fatigue of isolated mouse muscle due to isometric tetani and tetani with high power output. *Q J Exp Physiol* 74: 951–953, 1989.
  103. **Curtin NA, Edman KA.** Force-velocity relation for frog muscle fibres: effects of moderate fatigue and of intracellular acidification. *J Physiol* 475: 483–494, 1994.
  104. **Curtis BA.** Ca fluxes in single twitch muscle fibers. *J Gen Physiol* 50: 255–267, 1966.
  105. **Dahlstedt AJ, Katz A, Tavi P, Westerblad H.** Creatine kinase injection restores contractile function in creatine-kinase-deficient mouse skeletal muscle fibres. *J Physiol* 547: 395–403, 2003.
  106. **Dahlstedt AJ, Katz A, Westerblad H.** Role of myoplasmic phosphate in contractile function of skeletal muscle studies on creatine kinase deficient mice. *J Physiol* 533: 379–388, 2001.
  107. **Dahlstedt AJ, Katz A, Wieringa B, Westerblad H.** Is creatine kinase responsible for fatigue? Studies of isolated skeletal muscle deficient in creatine kinase. *FASEB J* 14: 982–990, 2000.
  108. **Dahlstedt AJ, Westerblad H.** Inhibition of creatine kinase reduces the fatigue-induced decrease of tetanic  $[\text{Ca}^{2+}]_i$  in mouse skeletal muscle. *J Physiol* 533: 639–649, 2001.
  109. **Damiani E, Sacchetto R, Margreth A.** Phosphorylation of anchoring protein by calmodulin protein kinase associated to the sarcoplasmic reticulum of rabbit fast-twitch muscle. *Biochem Biophys Res Commun* 279: 181–189, 2000.
  110. **Damiani E, Sacchetto R, Margreth A.** Variation of phospholamban in slow-twitch muscle sarcoplasmic reticulum between mammalian species and a link to the substrate specificity of endogenous  $\text{Ca}^{2+}$ -

- calmodulin-dependent protein kinase. *Biochim Biophys Acta* 1464: 231–241, 2000.
111. **Danieli-Betto D, Germinario E, Esposito A, Biral D, Betto R.** Effects of fatigue on sarcoplasmic reticulum and myofibrillar properties of rat single muscle fibers. *J Appl Physiol* 89: 891–898, 2000.
  112. **Dantzig JA, Goldman YE, Millar NC, Lacktis J, Homsher E.** Reversal of the cross-bridge force-generating transition by photogeneration of phosphate in rabbit psoas muscle fibres. *J Physiol* 451: 247–278, 1992.
  113. **Darnley GM, Duke AM, Steele DS, Macfarlane NG.** Effects of reactive oxygen species on aspects of excitation-contraction coupling in chemically skinned rabbit diaphragm muscle fibres. *Exp Physiol* 86: 161–168, 2001.
  114. **Davies CTM, White MJ.** Muscle weakness following eccentric work in man. *Pflügers Arch* 392: 168–171, 1981.
  115. **Davies KJ, Quintanilha AT, Brooks GA, Packer L.** Free radicals and tissue damage produced by exercise. *Biochem Biophys Res Commun* 107: 1198–1205, 1982.
  116. **Davies NW, Standen NB, Stanfield PR.** The effect of intracellular pH on ATP-dependent potassium channels of frog skeletal muscle. *J Physiol* 445: 549–568, 1992.
  117. **Dawson MJ, Gadian DG, Wilkie DR.** Muscle fatigue investigated by phosphorus nuclear magnetic resonance. *Nature* 274: 861–866, 1978.
  118. **Dawson MJ, Gadian DG, Wilkie DR.** Mechanical relaxation rate and metabolism studied in fatiguing muscle by phosphorus nuclear magnetic resonance. *J Physiol* 299: 465–484, 1980.
  119. **De Paula Brotto M, Van Leyen SA, Brotto LS, Jin JP, Nosek CM, Nosek TM.** Hypoxia/fatigue-induced degradation of troponin I and troponin C: new insights into physiologic muscle fatigue. *Pflügers Arch* 442: 738–744, 2001.
  120. **De Ruiter CJ, De Haan A.** Temperature effect on the force/velocity relationship of the fresh and fatigued human adductor pollicis muscle. *Pflügers Arch* 440: 163–170, 2000.
  121. **De Ruiter CJ, Jones DA, Sargeant AJ, De Haan A.** Temperature effect on the rates of isometric force development and relaxation in the fresh and fatigued human adductor pollicis muscle. *Exp Physiol* 84: 1137–1150, 1999.
  122. **De Ruiter CJ, May AM, Van Engelen BG, Wevers RA, Steenbergen-Spanjers GC, De Haan A.** Muscle function during repetitive moderate-intensity muscle contractions in myoadenylate deaminase-deficient Dutch subjects. *Clin Sci* 102: 531–539, 2002.
  123. **Debold EP, Dave H, Fitts RH.** Fiber type and temperature dependence of inorganic phosphate: implications for fatigue. *Am J Physiol Cell Physiol* 287: C673–C681, 2004.
  124. **Debold EP, Romatowski J, Fitts RH.** The depressive effect of  $P_i$  on the force-pCa relationship in skinned single muscle fibers is temperature dependent. *Am J Physiol Cell Physiol* 290: C1041–C1050, 2006.
  125. **Delbono O, Meissner G.** Sarcoplasmic reticulum  $Ca^{2+}$  release in rat slow- and fast-twitch muscles. *J Membr Biol* 151: 123–130, 1996.
  126. **Diamanti-Kandarakis E, Konstantinopoulos PA, Papailiou J, Kandarakis SA, Andreopoulos A, Sykiotis GP.** Erythropoietin abuse and erythropoietin gene doping: detection strategies in the genomic era. *Sports Med* 35: 831–840, 2005.
  127. **Diaz PT, She ZW, Davis WB, Clanton TL.** Hydroxylation of salicylate by the in vitro diaphragm: evidence for hydroxyl radical production during fatigue. *J Appl Physiol* 75: 540–545, 1993.
  128. **Dimauro S, Bonilla E, Davidson M, Hirano M, Schon EA.** Mitochondria in neuromuscular disorders. *Biochim Biophys Acta* 1366: 199–210, 1998.
  129. **Donaldson PL, Beam KG.** Calcium currents in a fast-twitch skeletal muscle of the rat. *J Gen Physiol* 82: 449–468, 1983.
  130. **Donaldson SK, Hermansen L, Bolles L.** Differential, direct effects of  $H^+$  on  $Ca^{2+}$ -activated force of skinned fibers from the soleus, cardiac and adductor magnus muscles of rabbits. *Pflügers Arch* 376: 55–65, 1978.
  131. **Drust B, Rasmussen P, Mohr M, Nielsen B, Nybo L.** Elevations in core and muscle temperature impairs repeated sprint performance. *Acta Physiol Scand* 183: 181–190, 2005.
  132. **Duke AM, Steele DS.** Effects of creatine phosphate on  $Ca^{2+}$  regulation by the sarcoplasmic reticulum in mechanically skinned rat skeletal muscle fibres. *J Physiol* 517: 447–458, 1999.
  133. **Duke AM, Steele DS.** Characteristics of phosphate-induced  $Ca^{2+}$  efflux from the SR in mechanically skinned rat skeletal muscle fibers. *Am J Physiol Cell Physiol* 278: C126–C135, 2000.
  134. **Duke AM, Steele DS.** Interdependent effects of inorganic phosphate and creatine phosphate on sarcoplasmic reticulum  $Ca^{2+}$  regulation in mechanically skinned rat skeletal muscle. *J Physiol* 531: 729–742, 2001.
  135. **Duke AM, Steele DS.** Mechanisms of reduced SR  $Ca^{2+}$  release induced by inorganic phosphate in rat skeletal muscle fibers. *Am J Physiol Cell Physiol* 281: C418–C429, 2001.
  136. **Dulhunty AF.** The dependence of membrane potential on extracellular chloride concentration in mammalian skeletal muscle fibres. *J Physiol* 276: 67–82, 1978.
  137. **Dulhunty AF.** Distribution of potassium and chloride permeability over the surface and T-tubule membranes of mammalian skeletal muscle. *J Membr Biol* 45: 293–310, 1979.
  138. **Dulhunty AF.** Heterogeneity of T-tubule geometry in vertebrate skeletal muscle fibres. *J Muscle Res Cell Motil* 5: 333–347, 1984.
  139. **Dulhunty AF.** The voltage-activation of contraction in skeletal muscle. *Prog Biophys Mol Biol* 57: 181–223, 1992.
  140. **Dulhunty AF.** Excitation-contraction coupling from the 1950s into the new millennium. *Clin Exp Pharmacol Physiol* 33: 763–772, 2006.
  141. **Duncan CJ.** Role of calcium in triggering rapid ultrastructural damage in muscle: a study with chemically skinned fibres. *J Cell Sci* 87: 581–594, 1987.
  142. **Dutka TL, Cole L, Lamb GD.** Calcium phosphate precipitation in the sarcoplasmic reticulum reduces action potential-mediated  $Ca^{2+}$  release in mammalian skeletal muscle. *Am J Physiol Cell Physiol* 289: C1502–C1512, 2005.
  143. **Dutka TL, Lamb GD.** Effect of lactate on depolarization-induced  $Ca^{2+}$  release in mechanically skinned skeletal muscle fibers. *Am J Physiol Cell Physiol* 278: C517–C525, 2000.
  144. **Dutka TL, Lamb GD.** Effect of carnosine on excitation-contraction coupling in mechanically-skinned rat skeletal muscle. *J Muscle Res Cell Motil* 25: 203–213, 2004.
  145. **Dutka TL, Lamb GD.** Effect of low cytoplasmic [ATP] on excitation-contraction coupling in fast-twitch muscle fibres of the rat. *J Physiol* 560: 451–468, 2004.
  146. **Dutka TL, Lamb GD.**  $Na^+K^+$  pumps in the transverse tubular system of skeletal muscle fibers preferentially use ATP from glycolysis. *Am J Physiol Cell Physiol* 293: C967–C977, 2007.
  147. **Dutka TL, Lamb GD.** Transverse tubular system depolarization reduces tetanic force in rat skeletal muscle fibers by impairing action potential repriming. *Am J Physiol Cell Physiol* 292: C2112–C2121, 2007.
  148. **Duty S, Allen DG.** The distribution of intracellular calcium concentration in isolated single fibres of mouse skeletal muscle during fatiguing stimulation. *Pflügers Arch* 427: 102–109, 1994.
  149. **Eberstein A, Sandow A.** Fatigue mechanisms in muscle fibers. In: *The Effect of Use and Disuse on the Neuromuscular Functions*, edited by Gutman E, Hink P. Amsterdam: Elsevier, 1963, p. 515–526.
  150. **Edman KA.** The velocity of unloaded shortening and its relation to sarcomere length and isometric force in vertebrate muscle fibres. *J Physiol* 291: 143–159, 1979.
  151. **Edman KA, Hwang JC.** The force-velocity relationship in vertebrate muscle fibres at varied tonicity of the extracellular medium. *J Physiol* 269: 255–272, 1977.
  152. **Edman KAP, Lou F.** Myofibrillar failure versus failure of activation during repetitive stimulation of frog muscle fibres. *J Physiol* 457: 655–673, 1992.
  153. **Edwards RH, McDonnell MJ, Hill DK.** A thermistor probe for myothermal measurements in man. *J Appl Physiol* 36: 511–513, 1974.
  154. **Edwards RHT, Hill DK, Jones DA.** Metabolic changes associated with the slowing of relaxation in fatigued mouse muscle. *J Physiol* 251: 287–301, 1975.
  155. **Edwards RHT, Hill DK, Jones DA, Merton PA.** Fatigue of long duration in human skeletal muscle after exercise. *J Physiol* 272: 769–778, 1977.
  156. **Eklblom BT.** Blood boosting and sport. *Baillieres Best Pract Res Clin Endocrinol Metab* 14: 89–98, 2000.



157. **Enoka RM, Stuart DG.** Neurobiology of muscle fatigue. *J Appl Physiol* 72: 1631–1648, 1992.
158. **Essen B, Jansson E, Henriksson J, Taylor AW, Saltin B.** Metabolic characteristics of fibre types in human skeletal muscle. *Acta Physiol Scand* 95: 153–165, 1975.
159. **Everts ME, Andersen JP, Clausen T, Hansen O.** Quantitative determination of  $\text{Ca}^{2+}$ -dependent  $\text{Mg}^{2+}$ -ATPase from sarcoplasmic reticulum in muscle biopsies. *Biochem J* 260: 443–448, 1989.
160. **Fabiato A, Fabiato F.** Effects of pH on the myofilaments and the sarcoplasmic reticulum of skinned cells from cardiac and skeletal muscles. *J Physiol* 276: 233–255, 1978.
161. **Favero TG, Pesssah IN, Klug GA.** Prolonged exercise reduces  $\text{Ca}^{2+}$  release in rat skeletal muscle sarcoplasmic reticulum. *Pflügers Arch* 422: 472–475, 1993.
162. **Favero TG, Zable AC, Abramson JJ.** Hydrogen peroxide stimulates the  $\text{Ca}^{2+}$  release channel from skeletal muscle sarcoplasmic reticulum. *J Biol Chem* 270: 25557–25563, 1995.
163. **Favero TG, Zable AC, Colter D, Abramson JJ.** Lactate inhibits  $\text{Ca}^{2+}$ -activated  $\text{Ca}^{2+}$ -channel activity from skeletal muscle sarcoplasmic reticulum. *J Appl Physiol* 82: 447–452, 1997.
164. **Ferenczi MA, Goldman YE, Simmons RM.** The dependence of force and shortening velocity on substrate concentration in skinned muscle fibres from *Rana temporaria*. *J Physiol* 350: 519–543, 1984.
165. **Filatov GN, Pinter MJ, Rich MM.** Resting potential-dependent regulation of the voltage sensitivity of sodium channel gating in rat skeletal muscle in vivo. *J Gen Physiol* 126: 161–172, 2005.
166. **Fink R, Hase S, Lüttgau HC, Wettwer E.** The effect of cellular energy reserves and internal calcium ions on the potassium conductance in skeletal muscle of the frog. *J Physiol* 336: 211–228, 1983.
167. **Fink R, Lüttgau HC.** An evaluation of the membrane constants and the potassium conductance in metabolically exhausted muscle fibres. *J Physiol* 263: 215–238, 1976.
168. **Fitts RH.** Cellular mechanisms of muscle fatigue. *Physiol Rev* 74: 49–94, 1994.
169. **Fong PY, Turner PR, Denetclaw WF, Steinhardt RA.** Increased activity of calcium leak channels in myotubes of Duchenne human and mdx mouse origin. *Science* 250: 673–676, 1990.
170. **Fowles JR, Green HJ.** Coexistence of potentiation and low-frequency fatigue during voluntary exercise in human skeletal muscle. *Can J Physiol Pharmacol* 81: 1092–1100, 2003.
171. **Franke C, Hatt H, Iazzo PA, Lehmann-Horn F.** Characteristics of  $\text{Na}^+$  channels and  $\text{Cl}^-$  conductance in resealed muscle fibre segments from patients with myotonic dystrophy. *J Physiol* 425: 391–405, 1990.
172. **Franzini-Armstrong C, Jorgensen AO.** Structure and development of E-C coupling units in skeletal muscle. *Annu Rev Physiol* 56: 509–534, 1994.
173. **Franzini-Armstrong C, Kenney LJ, Varriano-Marston E.** The structure of calsequestrin in triads of vertebrate skeletal muscle: a deep-etch study. *J Cell Biol* 105: 49–56, 1987.
174. **Fruen BR, Mickelson JR, Shomer NH, Roghair TR, Louis CF.** Regulation of the sarcoplasmic reticulum ryanodine receptor by inorganic phosphate. *J Biol Chem* 269: 192–198, 1994.
175. **Fryer MW, Owen VJ, Lamb GD, Stephenson DG.** Effects of creatine phosphate and  $\text{P}_i$  on  $\text{Ca}^{2+}$  movements and tension development in rat skinned skeletal muscle fibres. *J Physiol* 482: 123–140, 1995.
176. **Fryer MW, Stephenson DG.** Total and sarcoplasmic reticulum calcium contents of skinned fibres from rat skeletal muscle. *J Physiol* 493: 357–370, 1996.
177. **Funk C, Clark A Jr, Connett RJ.** How phosphocreatine buffers cyclic changes in ATP demand in working muscle. *Adv Exp Med Biol* 248: 687–692, 1989.
178. **Gage PW, Eisenberg RS.** Capacitance of the surface and transverse tubular membrane of frog sartorius muscle fibers. *J Gen Physiol* 53: 265–278, 1969.
179. **Gandevia SC.** Spinal and supraspinal factors in human muscle fatigue. *Physiol Rev* 81: 1725–1789, 2001.
180. **Garcia MC, Gonzalez-Serratos H, Morgan JP, Perreault CL, Rozycka M.** Differential activation of myofibrils during fatigue in phasic skeletal muscle cells. *J Muscle Res Cell Motil* 12: 412–424, 1991.
181. **Garland SJ, Enoka RM, Serrano LP, Robinson GA.** Behavior of motor units in human biceps brachii during a submaximal fatiguing contraction. *J Appl Physiol* 76: 2411–2419, 1994.
182. **Gilchrist JSC, Wang KKW, Katz S, Belcastro AN.** Calcium-activated neutral protease effects upon skeletal muscle sarcoplasmic reticulum protein structure and calcium release. *J Biol Chem* 267: 20857–20865, 1992.
183. **Gissel H, Clausen T.** Excitation induced  $\text{Ca}^{2+}$  uptake in rat skeletal muscle. *Am J Physiol Regul Integr Comp Physiol* 276: R331–R339, 1999.
184. **Glynn IM.** Transport adenosine triphosphatase in electric organ. The relation between ion transport and oxidative phosphorylation. *J Physiol* 169: 452–465, 1963.
185. **Gott RE, Nosek TM.** Changes in the intracellular milieu with fatigue or hypoxia depress contraction of skinned rabbit skeletal and cardiac muscle. *J Physiol* 412: 155–180, 1989.
186. **Goll DE, Thompson VF, Li H, Wei W, Cong J.** The calpain system. *Physiol Rev* 83: 731–801, 2003.
187. **Gong MC, Arbogast S, Guo Z, Mathenia J, Su W, Reid MB.** Calcium-independent phospholipase  $\text{A}_2$  modulates cytosolic oxidant activity and contractile function in murine skeletal muscle cells. *J Appl Physiol* 100: 399–405, 2006.
188. **Gonzalez E, Delbono O.** Recovery from fatigue in fast and slow single intact skeletal muscle fibers from aging mouse. *Muscle Nerve* 24: 1219–1224, 2001.
189. **Gonzalez-Alonso J, Teller C, Andersen SL, Jensen FB, Hyldig T, Nielsen B.** Influence of body temperature on the development of fatigue during prolonged exercise in the heat. *J Appl Physiol* 86: 1032–1039, 1999.
190. **Gonzalez-Serratos H, Somlyo AV, McCellan G, Shuman H, Borrero LM, Somlyo AP.** Composition of vacuoles and sarcoplasmic reticulum in fatigued muscle: electron probe analysis. *Proc Natl Acad Sci USA* 75: 1329–1333, 1978.
191. **Goodman C, Blazev R, Stephenson G.** Glycogen content and contractile responsiveness to T-system depolarization in skinned muscle fibres of the rat. *Clin Exp Pharmacol Physiol* 32: 749–756, 2005.
192. **Gordon AM, Homsher E, Regnier M.** Regulation of contraction in striated muscle. *Physiol Rev* 80: 853–924, 2000.
193. **Grabowski W, Lobsiger EA, Lüttgau HC.** The effect of repetitive stimulation at low frequencies upon the electrical and mechanical activity of single muscle fibres. *Pflügers Arch* 334: 222–239, 1972.
194. **Greenhaff PL, Söderlund K, Ren JM, Hultman E.** Energy metabolism in single human muscle fibres during intermittent contraction with occluded circulation. *J Physiol* 460: 443–453, 1993.
195. **Guharay F, Sachs F.** Stretch-activated single ion channel currents in tissue-cultured embryonic chick skeletal muscle. *J Physiol* 352: 685–701, 1984.
196. **Halliwell B, Gutteridge JMC.** *Free Radicals in Biology and Medicine*. New York: Oxford Univ. Press, 1998.
197. **Han JW, Thieleczek R, Varsanyi M, Heilmeyer LMG.** Compartmentalized ATP synthesis in skeletal muscle triads. *Biochemistry* 31: 377–384, 1992.
198. **Hancock CR, Brault JJ, Wiseman RW, Terjung RL, Meyer RA.**  $^{31}\text{P}$ -NMR observation of free ADP during fatiguing, repetitive contractions of murine skeletal muscle lacking AK1. *Am J Physiol Cell Physiol* 288: C1298–C1304, 2005.
199. **Hancock CR, Janssen E, Terjung RL.** Skeletal muscle contractile performance and ADP accumulation in adenylate kinase-deficient mice. *Am J Physiol Cell Physiol* 288: C1287–C1297, 2005.
200. **Hansen AK, Clausen T, Nielsen OB.** Effects of lactic acid and catecholamines on contractility in fast-twitch muscles exposed to hyperkalemia. *Am J Physiol Cell Physiol* 289: C104–C112, 2005.
201. **Hargreaves M.** Skeletal muscle metabolism during exercise in humans. *Clin Exp Pharmacol Physiol* 27: 225–228, 2000.
202. **Harrington D, Coats AJ.** Skeletal muscle abnormalities and evidence for their role in symptom generation in chronic heart failure. *Eur Heart J* 18: 1865–1872, 1997.
203. **Helander I, Westerblad H, Katz A.** Effects of glucose on contractile function,  $[\text{Ca}^{2+}]_i$  and glycogen in isolated mouse skeletal muscle. *Am J Physiol Cell Physiol* 282: C1306–C1312, 2002.



204. Hermansen L, Hultman E, Saltin B. Muscle glycogen during prolonged severe exercise. *Acta Physiol Scand* 71: 129–139, 1967.
205. Hickie IB, Hooker AW, Hadzi-Pavlovic D, Bennett BK, Wilson AJ, Lloyd AR. Fatigue in selected primary care settings: socio-demographic and psychiatric correlates. *Med J Aust* 164: 585–588, 1996.
206. Hidalgo C, Gonzalez ME, Lagos R. Characterization of the  $\text{Ca}^{2+}$ - or  $\text{Mg}^{2+}$ -ATPase of transverse tubule membranes isolated from rabbit skeletal muscle. *J Biol Chem* 258: 13937–13945, 1983.
207. Hidalgo C, Sanchez G, Barrientos G, Aracena-Parks P. A transverse tubule NADPH oxidase activity stimulates calcium release from isolated triads via ryanodine receptor Type 1 S-Glutathionylation. *J Biol Chem* 281: 26473–26482, 2006.
208. Hill AV, Kupalov P. Anaerobic and aerobic activity in isolated muscle. *Proc R Soc Lond B Biol Sci* 105: 313–322, 1929.
209. Hill CA, Harris RC, Kim HJ, Harris BD, Sale C, Boobis LH, Kim CK, Wise JA. Influence of beta-alanine supplementation on skeletal muscle carnosine concentrations and high intensity cycling capacity. *Amino Acids* 32: 225–233, 2007.
210. Hill CA, Thompson MW, Ruell PA, Thom JM, White MJ. Sarcoplasmic reticulum function and muscle contractile character following fatiguing exercise in humans. *J Physiol* 531: 871–878, 2001.
211. Hirata Y, Brotto M, Weisleder N, Chu Y, Lin P, Zhao X, Thornton A, Komazaki S, Takeshima H, Ma J, Pan Z. Uncoupling store-operated  $\text{Ca}^{2+}$  entry and altered  $\text{Ca}^{2+}$  release from sarcoplasmic reticulum through silencing of junctophilin genes. *Biophys J* 90: 4418–4427, 2006.
212. Hodgkin AL, Horowicz P. Movements of Na and K in single muscle fibres. *J Physiol* 145: 405–432, 1959.
213. Hodgkin AL, Horowicz P. The influence of potassium and chloride ions on the membrane potential of single muscle fibres. *J Physiol* 148: 127–160, 1959.
214. Hogan MC, Gladden LB, Kurdak SS, Poole DC. Increased [lactate] in working dog muscle reduces tension development independent of pH. *Med Sci Sports Exercise* 27: 371–377, 1995.
215. Hogan MC, Richardson RS, Haseler LJ. Human muscle performance and PCr hydrolysis with varied inspired oxygen fractions: a  $^{31}\text{P}$ -MRS study. *J Appl Physiol* 86: 1367–1373, 1999.
216. Hopf FW, Reddy P, Hong J, Steinhart RA. A capacitive calcium current in cultured skeletal muscle cells is mediated by the calcium-specific leak channel and inhibited by dihydropyridine compounds. *J Biol Chem* 271: 22358–22367, 1996.
217. Hoskins BK, Lipscomb S, Mulligan IP, Ashley CC. How do skinned skeletal muscle fibers relax? *Biochem Biophys Res Commun* 254: 330–333, 1999.
218. Howlett RA, Kelley KM, Grassi B, Gladden LB, Hogan MC. Caffeine administration results in greater tension development in previously fatigued canine muscle in situ. *Exp Physiol* 90: 873–879, 2005.
219. Howlett RA, Stary CM, Hogan MC. Recovery of force during postcontractile depression in single *Xenopus* muscle fibers. *Am J Physiol Regul Integr Comp Physiol* 280: R1469–R1475, 2001.
220. Hurne AM, O'Brien JJ, Wingrove D, Cherednichenko G, Allen PD, Beam KG, Pessah IN. Ryanodine receptor type 1 (RyR1) mutations C4958S and C4961S reveal excitation-coupled calcium entry (ECCE) is independent of sarcoplasmic reticulum store depletion. *J Biol Chem* 280: 36994–37004, 2005.
221. Inashima S, Matsunaga S, Yasuda T, Wada M. Effect of endurance training and acute exercise on sarcoplasmic reticulum function in rat fast- and slow-twitch skeletal muscles. *Eur J Appl Physiol* 89: 142–149, 2003.
222. Inesi G, De Meis L. Regulation of steady state filling in sarcoplasmic reticulum: roles of back-inhibition, leakage, slippage of the calcium pump. *J Biol Chem* 264: 5929–5936, 1989.
223. Isaeva EV, Shirokova N. Metabolic regulation of  $\text{Ca}^{2+}$  release in permeabilized mammalian skeletal muscle fibres. *J Physiol* 547: 453–462, 2003.
224. Jaimovich E, Venosa RA, Shrager P, Horowicz P. Density and distribution of tetrodotoxin receptors in normal and detubulated frog sartorius muscle. *J Gen Physiol* 67: 399–416, 1976.
225. Jansson E, Dudley GA, Norman B, Tesch PA. ATP and IMP in single human muscle fibres after high intensity exercise. *Clin Physiol* 7: 337–345, 1987.
226. Javesghani D, Magder SA, Barreiro E, Quinn MT, Hussain SN. Molecular characterization of a superoxide-generating NAD(P)H oxidase in the ventilatory muscles. *Am J Respir Crit Care Med* 165: 412–418, 2002.
227. Jones DA, Bigland-Ritchie B, Edwards RHT. Excitation frequency and muscle fatigue: mechanical responses during voluntary and stimulated contractions. *Exp Neurol* 64: 414–427, 1979.
228. Jones DA, De Ruiter J, De Haan A. Change in contractile properties of human muscle in relationship to the loss of power and slowing of relaxation seen with fatigue. *J Physiol* 576: 913–922, 2006.
229. Jones DA, Howell S, Roussos C, Edwards RHT. Low-frequency fatigue in isolated skeletal muscles and the effects of methylxanthines. *Clin Sci* 63: 161–167, 1982.
230. Juel C. Potassium and sodium shifts during in vitro isometric muscle contraction, the time course of the ion-gradient recovery. *Pflügers Arch* 406: 458–463, 1986.
231. Juel C. Muscle action potential propagation velocity changes during activity. *Muscle Nerve* 11: 714–719, 1988.
232. Juel C, Pilegaard H, Nielsen JJ, Bangsbo J. Interstitial  $\text{K}^{+}$  in human skeletal muscle during and after dynamic graded exercise determined by microdialysis. *Am J Physiol Regul Integr Comp Physiol* 278: R400–R406, 2000.
233. Jurkat-Rott K, Fauler M, Lehmann-Horn F. Ion channels and ion transporters of the transverse tubular system of skeletal muscle. *J Muscle Res Cell Motil* 27: 275–290, 2006.
234. Kabbara AA, Allen DG. The role of calcium stores in fatigue of isolated single muscle fibres from the cane toad. *J Physiol* 519: 169–176, 1999.
235. Kabbara AA, Allen DG. The use of fluo-5N to measure sarcoplasmic reticulum calcium in single muscle fibres of the cane toad. *J Physiol* 534: 87–97, 2001.
236. Kabbara AA, Nguyen LT, Stephenson GMM, Allen DG. Intracellular calcium during fatigue of cane toad skeletal muscle in the absence of glucose. *J Muscle Res Cell Motil* 21: 481–489, 2000.
237. Kabbara AA, Stephenson DG. Effects of  $\text{Mg}^{2+}$  on  $\text{Ca}^{2+}$  handling by the sarcoplasmic reticulum in skinned skeletal and cardiac muscle fibres. *Pflügers Arch* 428: 331–339, 1994.
238. Karatzafieri C, De Haan A, Ferguson RA, Van Mechelen W, Sargeant AJ. Phosphocreatine and ATP content in human single muscle fibres before and after maximum dynamic exercise. *Pflügers Arch* 442: 467–474, 2001.
239. Karelis AD, Marcil M, Peronnet F, Gardiner PF. Effect of lactate infusion on M-wave characteristics and force in the rat plantaris muscle during repeated stimulation in situ. *J Appl Physiol* 96: 2133–2138, 2004.
240. Karlsson J, Funderburk CF, Essen B, Lind AR. Constituents of human muscle in isometric fatigue. *J Appl Physiol* 38: 208–211, 1975.
241. Kasuga N, Umazume Y. Deterioration induced by physiological concentration of calcium ions in skinned muscle fibres. *J Muscle Res Cell Motil* 11: 41–47, 1990.
242. Kim KC, Caswell AH, Brunschwig JP, Brandt NR. Identification of a new subpopulation of triad junctions isolated from skeletal muscle; morphological correlations with intact muscle. *J Membr Biol* 113: 221–235, 1990.
243. Kindig CA, Howlett RA, Stary CM, Walsh B, Hogan MC. Effects of acute creatine kinase inhibition on metabolism and tension development in isolated single myocytes. *J Appl Physiol* 98: 541–549, 2005.
244. Knuth S, Dave H, Peters JR, Fitts RH. Low cell pH depresses peak power in rat skeletal muscle fibres at both 30 and 15 degrees centigrade: Implications for muscle fatigue. *J Physiol* 575: 887–899, 2006.
245. Kolbeck RC, She ZW, Callahan LA, Nosek TM. Increased superoxide production during fatigue in the perfused rat diaphragm. *Am J Respir Crit Care Med* 156: 140–145, 1997.
246. Konishi M, Suda N, Kurihara S. Fluorescence signals from the  $\text{Mg}^{2+}/\text{Ca}^{2+}$  indicator fura-2 in frog skeletal muscle fibers. *Biophys J* 64: 223–239, 1993.

247. **Korge P, Campbell KB.** The importance of ATPase microenvironment in muscle fatigue: a hypothesis. *Int J Sports Med* 16: 172–179, 1995.
248. **Kristensen M, Albertsen J, Rentsch M, Juel C.** Lactate and force production in skeletal muscle. *J Physiol* 562: 521–526, 2005.
249. **Kristensen M, Hansen T, Juel C.** Membrane proteins involved in potassium shifts during muscle activity and fatigue. *Am J Physiol Regul Integr Comp Physiol* 290: R766–R772, 2006.
250. **Kugelberg E, Lindegren B.** Transmission and contraction fatigue of rat motor units in relation to succinate dehydrogenase activity of motor unit fibres. *J Physiol* 288: 285–300, 1979.
251. **Kurebayashi N, Ogawa Y.** Depletion of  $\text{Ca}^{2+}$  in the sarcoplasmic reticulum stimulates  $\text{Ca}^{2+}$  entry into mouse skeletal muscle fibres. *J Physiol* 533: 185–199, 2001.
252. **Lamb GD.** DHP receptors and excitation-contraction coupling. *J Muscle Res Cell Motil* 13: 394–405, 1992.
253. **Lamb GD.** Rippling muscle disease may be caused by “silent” action potentials in the tubular system of skeletal muscle fibers. *Muscle Nerve* 31: 652–658, 2005.
254. **Lamb GD, Junankar PR, Stephenson DG.** Raised intracellular  $[\text{Ca}^{2+}]$  abolishes excitation-contraction coupling in skeletal muscle fibres of rat and toad. *J Physiol* 489: 349–362, 1995.
255. **Lamb GD, Posterino GS.** Effects of oxidation and reduction on contractile function in skeletal muscle fibres of the rat. *J Physiol* 546: 149–163, 2003.
256. **Lamb GD, Recupero E, Stephenson DG.** Effect of myoplasmic pH on excitation-contraction coupling in skeletal muscle fibres of the toad. *J Physiol* 448: 211–224, 1992.
257. **Lamb GD, Stephenson DG.** Effects of  $\text{Mg}^{2+}$  on the control of  $\text{Ca}^{2+}$  release in skeletal muscle fibres of the toad. *J Physiol* 434: 507–528, 1991.
258. **Lamb GD, Stephenson DG.** Effects of intracellular pH and  $[\text{Mg}^{2+}]$  on excitation-contraction coupling in skeletal muscle fibres of the rat. *J Physiol* 478: 331–339, 1994.
259. **Lamb GD, Stephenson DG, Stienen GJ.** Effects of osmolality and ionic strength on the mechanism of  $\text{Ca}^{2+}$  release in skinned skeletal muscle fibres of the toad. *J Physiol* 464: 629–648, 1993.
260. **Lamb GD, Walsh T.** Calcium currents, charge movement and dihydropyridine binding in fast- and slow-twitch muscles of rat and rabbit. *J Physiol* 393: 595–617, 1987.
261. **Lamont C, Miller DJ.** Calcium sensitizing action of carnosine and other endogenous imidazoles in chemically skinned striated muscle. *J Physiol* 454: 421–434, 1992.
262. **Lännergren J, Bruton JD, Westerblad H.** Vacuole formation in fatigued single muscle fibres from frog and mouse. *J Muscle Res Cell Motil* 20: 19–32, 1999.
263. **Lännergren J, Bruton JD, Westerblad H.** Vacuole formation in fatigued skeletal muscle fibres from frog and mouse: effects of extracellular lactate. *J Physiol* 526: 597–611, 2000.
264. **Lännergren J, Larsson L, Westerblad H.** A novel type of delayed tension reduction observed in rat motor units after intense activity. *J Physiol* 412: 267–276, 1989.
265. **Lännergren J, Westerblad H.** Force and membrane potential during and after fatiguing, continuous high-frequency stimulation of single *Xenopus* muscle fibres. *Acta Physiol Scand* 128: 359–368, 1986.
266. **Lännergren J, Westerblad H.** Action potential fatigue in single skeletal muscle fibres of *Xenopus*. *Acta Physiol Scand* 129: 311–318, 1987.
267. **Lännergren J, Westerblad H.** The temperature dependence of isometric contractions of single, intact fibres dissected from a mouse foot muscle. *J Physiol* 390: 285–293, 1987.
268. **Lännergren J, Westerblad H.** Force decline due to fatigue and intracellular acidification in isolated fibres from mouse skeletal muscle. *J Physiol* 434: 307–322, 1991.
269. **Lännergren J, Westerblad H, Bruton JD.** Slow recovery of force in single skeletal muscle fibres. *Acta Physiol Scand* 156: 193–202, 1996.
270. **Lännergren J, Westerblad H, Bruton JD.** Changes in mitochondrial  $\text{Ca}^{2+}$  detected with Rhod-2 in single frog and mouse skeletal muscle fibres during and after repeated tetanic contractions. *J Muscle Res Cell Motil* 22: 265–275, 2001.
271. **Lännergren J, Westerblad H, Bruton JD.** Dynamic vacuolation in skeletal muscle fibres after fatigue. *Cell Biol Int* 26: 911–920, 2002.
272. **Launikonis BS, Barnes M, Stephenson DG.** Identification of the coupling between skeletal muscle store-operated  $\text{Ca}^{2+}$  entry and the inositol trisphosphate receptor. *Proc Natl Acad Sci USA* 100: 2941–2944, 2003.
273. **Launikonis BS, Stephenson DG.** Osmotic properties of the sealed tubular system of toad and rat skeletal muscle. *J Gen Physiol* 123: 231–247, 2004.
274. **Launikonis BS, Zhou J, Royer L, Shannon TR, Brum G, Rios E.** Confocal imaging of  $[\text{Ca}^{2+}]$  in cellular organelles by SEER, shifted excitation and emission ratioing of fluorescence. *J Physiol* 567: 523–543, 2005.
275. **Launikonis BS, Zhou J, Royer L, Shannon TR, Brum G, Rios E.** Depletion “skraps” and dynamic buffering inside the cellular calcium store. *Proc Natl Acad Sci USA* 103: 2982–2987, 2006.
276. **Lauritzen HP, Ploug T, Prats C, Tavares JM, Galbo H.** Imaging of insulin signaling in skeletal muscle of living mice shows major role of T-tubules. *Diabetes* 55: 1300–1306, 2006.
277. **Laver DR, Eager KR, Taoube L, Lamb GD.** Effects of cytoplasmic and luminal pH on  $\text{Ca}^{2+}$  release channels from rabbit skeletal muscle. *Biophys J* 78: 1835–1851, 2000.
278. **Laver DR, Lenz GK, Lamb GD.** Regulation of the calcium release channel from rabbit skeletal muscle by the nucleotides ATP, AMP, IMP and adenosine. *J Physiol* 537: 763–778, 2001.
279. **Laver DR, Lenz GKE, Dulhunty AF.** Phosphate ion channels in the sarcoplasmic reticulum of rabbit skeletal muscle. *J Physiol* 537: 763–778, 2001.
280. **Laver DR, O'Neill ER, Lamb GD.** Luminal  $\text{Ca}^{2+}$ -regulated  $\text{Mg}^{2+}$  inhibition of skeletal RyRs reconstituted as isolated channels or coupled clusters. *J Gen Physiol* 124: 741–758, 2004.
281. **Laver DR, Owen VJ, Junankar PR, Taske NL, Dulhunty AF, Lamb GD.** Reduced inhibitory effect of  $\text{Mg}^{2+}$  on ryanodine receptor- $\text{Ca}^{2+}$  release channels in malignant hyperthermia. *Biophys J* 73: 1913–1924, 1997.
282. **Leberer E, Pette D.** Immunochemical quantification of sarcoplasmic reticulum Ca-ATPase, of calsequestrin and of parvalbumin in rabbit skeletal muscles of defined fiber composition. *Eur J Biochem* 156: 489–496, 1986.
283. **Lee JA, Westerblad H, Allen DG.** Changes in tetanic and resting  $[\text{Ca}^{2+}]$  during fatigue and recovery of single muscle fibres from *Xenopus laevis*. *J Physiol* 433: 307–326, 1991.
284. **Leppik JA, Aughey RJ, Medved I, Fairweather I, Carey MF, McKenna MJ.** Prolonged exercise to fatigue in humans impairs skeletal muscle  $\text{Na}^{+}$ - $\text{K}^{+}$ -ATPase activity, sarcoplasmic reticulum  $\text{Ca}^{2+}$  release,  $\text{Ca}^{2+}$  uptake. *J Appl Physiol* 97: 1414–1423, 2004.
285. **Li JL, Wang XN, Fraser SF, Carey MF, Wrigley TV, McKenna MJ.** Effects of fatigue and training on sarcoplasmic reticulum  $\text{Ca}^{2+}$  regulation in human skeletal muscle. *J Appl Physiol* 92: 912–922, 2002.
286. **Lindinger MI, Heigenhauser GJ, Spriet LL.** Effects of intense swimming and tetanic electrical stimulation on skeletal muscle ions and metabolites. *J Appl Physiol* 63: 2331–2339, 1987.
287. **Lunde PK, Dahlstedt AJ, Bruton JD, Lännergren J, Thoren P, Sejersted OM, Westerblad H.** Contraction and intracellular  $\text{Ca}^{2+}$  handling in isolated skeletal muscle of rats with congestive heart failure. *Circ Res* 88: 1299–1305, 2001.
288. **Lunde PK, Sejersted OM, Schiøtz Thorud HM, Tønnessen T, Henriksen UL, Christensen G, Westerblad H, Bruton J.** Effects of congestive heart failure on  $\text{Ca}^{2+}$  handling in skeletal muscle during fatigue. *Circ Res* 98: 1514–1519, 2006.
289. **Lunde PK, Sjaastad I, Schiøtz Thorud HM, Sejersted OM.** Skeletal muscle disorders in heart failure. *Acta Physiol Scand* 171: 277–294, 2001.
290. **Lüttgau HC, Spiecker W.** The effects of calcium deprivation upon mechanical and electrophysiological parameters in skeletal muscle fibres of the frog. *J Physiol* 296: 411–429, 1979.
291. **Lynch GS, Schertzer JD, Ryall JG.** Therapeutic approaches for muscle wasting disorders. *Pharmacol Ther.* In press.
292. **Lynch RE, Fridovich I.** Permeation of the erythrocyte stroma by superoxide radical. *J Biol Chem* 253: 4697–4699, 1978.



293. **Lytton J, Westlin M, Burk SE, Shull GE, MacLennan DH.** Functional comparisons between isoforms of the sarcoplasmic or endoplasmic reticulum family of calcium pumps. *J Biol Chem* 267: 14483–14489, 1992.
294. **Ma J, Fill M, Knudson CM, Campbell KP, Coronado R.** Ryanodine receptor of skeletal muscle is a gap junction-type channel. *Science* 242: 99–102, 1988.
295. **MacDonald WA, Stephenson DG.** Effects of ADP on sarcoplasmic reticulum function in mechanically skinned skeletal muscle fibres of the rat. *J Physiol* 532: 499–508, 2001.
296. **MacDonald WA, Stephenson DG.** Effect of ADP on slow-twitch muscle fibres of the rat: implications for muscle fatigue. *J Physiol* 573: 187–198, 2006.
297. **Madsen K, Ertbjerg P, Djurhuus MS, Pedersen PK.** Calcium content and respiratory control index of skeletal muscle mitochondria during exercise and recovery. *Am J Physiol Endocrinol Metab* 271: E1044–E1050, 1996.
298. **Manning DR, Stull JT.** Myosin light chain phosphorylation-dephosphorylation in mammalian skeletal muscle. *Am J Physiol Cell Physiol* 242: C234–C241, 1982.
299. **Margreth A, Damiani E, Tobaldin G.** Ratio of dihydropyridine to ryanodine receptors in mammalian and frog twitch muscles in relation to the mechanical hypothesis of excitation-contraction coupling. *Biochem Biophys Res Commun* 197: 1303–1311, 1993.
300. **Marsden CD, Meadows JC, Merton PA.** Isolated single motor units in human muscle and their rate of discharge during maximal voluntary effort. *J Physiol* 217: 12P–13P, 1971.
301. **Martyn DA, Gordon AM.** Force and stiffness in glycerinated rabbit psoas fibers. Effects of calcium and elevated phosphate. *J Gen Physiol* 99: 795–816, 1992.
302. **McFadden LK, McComas AJ.** Late depression of muscle excitability in humans after fatiguing stimulation. *J Physiol* 496: 851–855, 1996.
303. **McKenna MJ, Medved I, Goodman CA, Brown MJ, Bjorksten AR, Murphy KT, Petersen AC, Sostaric S, Gong X.** N-Acetylcysteine attenuates the decline in muscle  $\text{Na}^+$ ,  $\text{K}^+$ -pump activity and delays fatigue during prolonged exercise in humans. *J Physiol* 576: 279–288, 2006.
304. **Medved I, Brown MJ, Bjorksten AR, McKenna MJ.** Effects of intravenous N-acetylcysteine infusion on time to fatigue and potassium regulation during prolonged cycling exercise. *J Appl Physiol* 96: 211–217, 2004.
305. **Meissner G, Darling E, Eveleth J.** Kinetics of rapid  $\text{Ca}^{2+}$  release by sarcoplasmic reticulum. Effects of  $\text{Ca}^{2+}$ ,  $\text{Mg}^{2+}$ , adenine nucleotides. *Biochemistry* 25: 236–244, 1986.
306. **Melzer W, Herrmann-Frank A, Lüttgau HC.** The role of  $\text{Ca}^{2+}$  ions in excitation-contraction coupling of skeletal muscle fibres. *Biochim Biophys Acta* 1241: 59–116, 1995.
307. **Melzer W, Rios E, Schneider MF.** Time course of calcium release and removal in skeletal muscle fibers. *Biophys J* 45: 637–641, 1984.
308. **Merton PA.** Voluntary strength and fatigue. *J Physiol* 123: 553–564, 1954.
309. **Metzger JM.** Effects of phosphate and ADP on shortening velocity during maximal and submaximal calcium activation of the thin filament in skeletal muscle fibers. *Biophys J* 70: 409–417, 1996.
310. **Metzger JM, Fitts RH.** Role of intracellular pH in muscle fatigue. *J Appl Physiol* 62: 1392–1397, 1987.
311. **Meyer RA, Sweeney HL, Kushmerick MJ.** A simple analysis of the “phosphocreatine shuttle.” *Am J Physiol Cell Physiol* 246: C365–C377, 1984.
312. **Meyer RA, Terjung RL.** Differences in ammonia and adenylate metabolism in contracting fast and slow muscle. *Am J Physiol Cell Physiol* 237: C111–C118, 1979.
313. **Mikkelsen UR, Fredsted A, Gissel H, Clausen T.** Excitation-induced  $\text{Ca}^{2+}$  influx and muscle damage in the rat: loss of membrane integrity and impaired force recovery. *J Physiol* 559: 271–285, 2004.
314. **Millar NC, Homsher E.** The effect of phosphate and calcium on force generation in glycerinated rabbit skeletal muscle fibers: a steady-state and transient kinetic study. *J Biol Chem* 265: 20234–20240, 1990.
315. **Moopanar TR, Allen DG.** Reactive oxygen species reduce myofibrillar  $\text{Ca}^{2+}$ -sensitivity in fatiguing mouse skeletal muscle at 37°C. *J Physiol* 564: 189–199, 2005.
316. **Moopanar TR, Allen DG.** The activity-induced reduction of myofibrillar  $\text{Ca}^{2+}$  sensitivity in mouse skeletal muscle is reversed by dithiothreitol. *J Physiol* 571: 191–200, 2006.
317. **Moore RL, Stull JT.** Myosin light chain phosphorylation in fast and slow skeletal muscles in situ. *Am J Physiol Cell Physiol* 247: C462–C471, 1984.
318. **Mosso A.** *Fatigue*. London: Swan Sonnenschein, 1904.
319. **Mulligan IP, Palmer RE, Lipscomb S, Hoskins B, Ashley CC.** The effect of phosphate on the relaxation of frog skeletal muscle. *Pflügers Arch* 437: 393–399, 1999.
320. **Murphy RM, Snow RJ, Lamb GD.**  $\mu$ -Calpain and calpain-3 are not autolyzed with exhaustive exercise in humans. *Am J Physiol Cell Physiol* 290: C116–C122, 2006.
321. **Murphy RM, Stephenson DG, Lamb GD.** Effect of creatine on contractile force and sensitivity in mechanically skinned single fibers from rat skeletal muscle. *Am J Physiol Cell Physiol* 287: C1589–C1595, 2004.
322. **Murphy RM, Verburg E, Lamb GD.**  $\text{Ca}^{2+}$ -activation of diffusible and bound pools of  $\mu$ -calpain in rat skeletal muscle. *J Physiol* 576: 595–612, 2006.
323. **Nagesser AS, Van Der Laarse WJ, Elzinga G.** Metabolic changes with fatigue in different types of single muscle fibres of *Xenopus laevis*. *J Physiol* 448: 511–523, 1992.
324. **Nagesser AS, Van Der Laarse WJ, Elzinga G.** ATP formation and ATP hydrolysis during fatiguing, intermittent stimulation of different types of single muscle fibres from *Xenopus laevis*. *J Muscle Res Cell Motil* 14: 608–618, 1993.
325. **Nakajima S, Gilai A.** Radial propagation of muscle action potential along the tubular system examined by potential-sensitive dyes. *J Gen Physiol* 76: 751–762, 1980.
326. **Nakajima S, Nakajima Y, Peachey LD.** Speed of repolarization and morphology of glycerol-treated frog muscle fibres. *J Physiol* 234: 465–480, 1973.
327. **Nakamura J, Tajima G, Sato C, Furukohri T, Konishi K.** Substrate regulation of calcium binding in  $\text{Ca}^{2+}$ -ATPase molecules of the sarcoplasmic reticulum. I. Effect of ATP. *J Biol Chem* 277: 24180–24190, 2002.
328. **Needham DM.** *Machina Carnis* Cambridge, UK: Cambridge Univ. Press, 1971.
329. **Nethery D, Callahan LA, Stofan D, Mattera R, Dimarco A, Supinski G.**  $\text{PLA}_2$  dependence of diaphragm mitochondrial formation of reactive oxygen species. *J Appl Physiol* 89: 72–80, 2000.
330. **Nicoletti I, Ciccoira M, Zanolla L, Franceschini L, Brighetti G, Pilati M, Zardini P.** Skeletal muscle abnormalities in chronic heart failure patients: relation to exercise capacity and therapeutic implications. *Congest Heart Fail* 9: 148–154, 2003.
331. **Nielsen JJ, Kristensen M, Hellsten Y, Bangsbo J, Juel C.** Localization and function of ATP-sensitive potassium channels in human skeletal muscle. *Am J Physiol Regul Integr Comp Physiol* 284: R558–R563, 2003.
332. **Nielsen JJ, Mohr M, Klarskov C, Kristensen M, Krstrup P, Juel C, Bangsbo J.** Effects of high-intensity intermittent training on potassium kinetics and performance in human skeletal muscle. *J Physiol* 554: 857–870, 2004.
333. **Nielsen OB, De Paoli F, Overgaard K.** Protective effects of lactic acid on force production in rat skeletal muscle. *J Physiol* 536: 161–166, 2001.
334. **Nielsen OB, MacDonald W, Stephenson DG, Clausen T.** Distribution of  $\text{Na}^+$ - $\text{K}^+$  pumps in skeletal muscle and its significance for maintenance of T-tubular  $\text{K}^+$  homeostasis. *Proc Physiol Soc* 4: C4, 2006.
335. **Nielsen OB, Ørtenblad N, Lamb GD, Stephenson DG.** Excitability of the T-tubular system in rat skeletal muscle: roles of  $\text{K}^+$  and  $\text{Na}^+$  gradients and  $\text{Na}^+$ - $\text{K}^+$  pump activity. *J Physiol* 557: 133–146, 2004.
336. **Norman B, Sabina RL, Jansson E.** Regulation of skeletal muscle ATP catabolism by AMPD1 genotype during sprint exercise in asymptomatic subjects. *J Appl Physiol* 91: 258–264, 2001.
337. **Okamoto K, Wang W, Rounds J, Chambers EA, Jacobs DO.** ATP from glycolysis is required for normal sodium homeostasis in



- resting fast-twitch rodent skeletal muscle. *Am J Physiol Endocrinol Metab* 281: E479–E488, 2001.
338. **Overgaard K, Fredsted A, Hyldal A, Ingemann-Hansen T, Gissel H, Clausen T.** Effects of running distance and training on  $\text{Ca}^{2+}$  content and damage in human muscle. *Med Sci Sports Exercise* 36: 821–829, 2004.
  339. **Overgaard K, Nielsen OB.** Activity-induced recovery of excitability in  $\text{K}^{+}$ -depressed rat soleus muscle. *Am J Physiol Regul Integr Comp Physiol* 280: R48–R55, 2001.
  340. **Overgaard K, Nielsen OB, Clausen T.** Effects of reduced electrochemical  $\text{Na}^{+}$  gradient on contractility in skeletal muscle: role of the  $\text{Na}^{+}$ - $\text{K}^{+}$  pump. *Pflügers Arch* 434: 457–465, 1997.
  341. **Overgaard K, Nielsen OB, Flatman JA, Clausen T.** Relations between excitability and contractility in rat soleus muscle: role of the  $\text{Na}^{+}$ - $\text{K}^{+}$  pump and  $\text{Na}^{+}/\text{K}^{+}$  gradients. *J Physiol* 518: 215–225, 1999.
  342. **Owen VJ, Lamb GD, Stephenson DG.** Effect of low [ATP] on depolarization-induced  $\text{Ca}^{2+}$  release in skeletal muscle fibres of the toad. *J Physiol* 493: 309–315, 1996.
  343. **Owen VJ, Lamb GD, Stephenson DG, Fryer MW.** Relationship between depolarization-induced force response and  $\text{Ca}^{2+}$  content in skeletal muscle fibres of rat and toad. *J Physiol* 498: 571–586, 1997.
  344. **Pan Z, Yang D, Nagaraj RY, Nosek TA, Nishi M, Takeshima H, Cheng H, Ma J.** Dysfunction of store-operated calcium channel in muscle cells lacking  $\text{mg29}$ . *Nat Cell Biol* 4: 379–383, 2002.
  345. **Pape PC, Jong DS, Chandler WK.** Effects of partial sarcoplasmic reticulum calcium depletion on calcium release in frog cut muscle fibers equilibrated with 20 mM EGTA. *J Gen Physiol* 112: 263–295, 1998.
  346. **Pape PC, Jong DS, Chandler WK, Baylor SM.** Effect of fura-2 on action potential-stimulated calcium release in cut twitch fibers from frog muscle. *J Gen Physiol* 102: 295–332, 1993.
  347. **Pate E, Bhimani M, Franks-Skiba K, Cooke R.** Reduced effect of pH on skinned rabbit psoas muscle mechanics at high temperatures: implications for fatigue. *J Physiol* 486: 689–694, 1995.
  348. **Pate E, Cooke R.** Addition of phosphate to active muscle fibers probes actomyosin states within the powerstroke. *Pflügers Arch* 414: 73–81, 1989.
  349. **Pathare N, Walter GA, Stevens JE, Yang Z, Okerke E, Gibbs JD, Esterhai JL, Scarborough MT, Gibbs CP, Sweeney HL, Vandenborne K.** Changes in inorganic phosphate and force production in human skeletal muscle after cast immobilization. *J Appl Physiol* 98: 307–314, 2005.
  350. **Pedersen TH, Clausen T, Nielsen OB.** Loss of force induced by high extracellular  $[\text{K}^{+}]$  in rat muscle: effect of temperature, lactic acid and beta2-agonist. *J Physiol* 551: 277–286, 2003.
  351. **Pedersen TH, De Paoli F, Nielsen OB.** Increased excitability of acidified skeletal muscle: role of chloride conductance. *J Gen Physiol* 125: 237–246, 2005.
  352. **Pedersen TH, Nielsen OB, Lamb GD, Stephenson DG.** Intracellular acidosis enhances the excitability of the working muscle. *Science* 305: 1144–1147, 2004.
  353. **Persechini A, Stull JT, Cooke R.** The effect of myosin phosphorylation on the contractile properties of skinned rabbit skeletal muscle fibers. *J Biol Chem* 260: 7951–7954, 1985.
  354. **Pette D, Staron RS.** Myosin isoforms, muscle fiber types, transitions. *Microsc Res Tech* 50: 500–509, 2000.
  355. **Phillips SK, Wiseman RW, Woledge RC, Kushmerick MJ.** The effect of metabolic fuel on force production and resting inorganic phosphate levels in mouse skeletal muscle. *J Physiol* 462: 135–146, 1993.
  356. **Plant DR, Gregorevic P, Williams DA, Lynch GS.** Redox modulation of maximum force production of fast- and slow twitch skeletal muscles of rats and mice. *J Appl Physiol* 90: 832–838, 2001.
  357. **Posterino GS, Cellini MA, Lamb GD.** Effects of oxidation and cytosolic redox conditions on excitation-contraction coupling in rat skeletal muscle. *J Physiol* 547: 807–823, 2003.
  358. **Posterino GS, Dutka TL, Lamb GD.**  $\text{L}(+)$ -Lactate does not affect twitch and tetanic responses in mechanically skinned mammalian muscle fibres. *Pflügers Arch* 442: 197–203, 2001.
  359. **Posterino GS, Fryer MW.** Mechanisms underlying phosphate-induced failure of  $\text{Ca}^{2+}$  release in single skinned skeletal muscle fibres of the rat. *J Physiol* 512: 97–108, 1998.
  360. **Posterino GS, Lamb GD.** Effect of sarcoplasmic reticulum  $\text{Ca}^{2+}$  content on action potential-induced  $\text{Ca}^{2+}$  release in rat skeletal muscle fibres. *J Physiol* 551: 219–237, 2003.
  361. **Posterino GS, Lamb GD, Stephenson DG.** Twitch and tetanic force responses and longitudinal propagation of action potentials in skinned skeletal muscle fibres of the rat. *J Physiol* 527: 131–137, 2000.
  362. **Powers SK, Lennon SL.** Analysis of cellular responses to free radicals: focus on exercise and skeletal muscle. *Proc Nutr Soc* 58: 1025–1033, 1999.
  363. **Proske U, Morgan DL.** Muscle damage from eccentric exercise: mechanism, mechanical signs, adaptation and clinical applications. *J Physiol* 537: 333–345, 2001.
  364. **Radzyukevich T, Edman KA.** Effects of intracellular acidification and varied temperature on force, stiffness, speed of shortening in frog muscle fibers. *Am J Physiol Cell Physiol* 287: C106–C113, 2004.
  365. **Ranatunga KW.** Effects of acidosis on tension development in mammalian skeletal muscle. *Muscle Nerve* 10: 439–445, 1987.
  366. **Rassier DE, Macintosh BR.** Coexistence of potentiation and fatigue in skeletal muscle. *Braz J Med Biol Res* 33: 499–508, 2000.
  367. **Raynaud F, Bonnal C, Fernandez E, Bremaud L, Cerutti M, Lebart Mc Roustan C, Ouali A, Benyamin Y.** The calpain 1- $\alpha$ -actinin interaction. Resting complex between the calcium-dependent protease and its target in cytoskeleton. *Eur J Biochem* 270: 4662–4670, 2003.
  368. **Reid MB.** Invited review: redox modulation of skeletal muscle contraction: what we know and what we don't. *J Appl Physiol* 90: 724–731, 2001.
  369. **Reid MB, Durham WJ.** Generation of reactive oxygen and nitrogen species in contracting skeletal muscle: potential impact on aging. *Ann NY Acad Sci* 959: 108–116, 2002.
  370. **Reid MB, Stokic DS, Koch SM, Khawli FA, Leis AA.** N-acetylcysteine inhibits muscle fatigue in humans. *J Clin Invest* 94: 2468–2474, 1994.
  371. **Reid MR, Haack KE, Franchek KM, Valberg PA, Kobzik L, West MS.** Reactive oxygen in skeletal muscle I Intracellular oxidant kinetics and fatigue in vitro. *J Appl Physiol* 73: 1797–1804, 1992.
  372. **Reiken S, Gaburjakova M, Guatimosim S, Gomez AM, D'armiento J, Burkhoff D, Wang J, Vassort G, Lederer WJ, Marks AR.** Protein kinase A phosphorylation of the cardiac calcium release channel (ryanodine receptor) in normal and failing hearts. Role of phosphatases and response to isoproterenol. *J Biol Chem* 278: 444–453, 2003.
  373. **Renaud JM, Allard Y, Mainwood GW.** Is the change in intracellular pH during fatigue large enough to be the main cause of fatigue? *Can J Physiol Pharmacol* 64: 764–767, 1986.
  374. **Renaud JM, Light P.** Effects of  $\text{K}^{+}$  on the twitch and tetanic contraction in the sartorius muscle of the frog, *Rana pipiens*. Implication for fatigue in vivo. *Can J Physiol Pharmacol* 70: 1236–1246, 1992.
  375. **Rich MM, Pinter MJ.** Crucial role of sodium channel fast inactivation in muscle fibre inexcitability in a rat model of critical illness myopathy. *J Physiol* 547: 555–566, 2003.
  376. **Richardson RS, Duteil S, Wary C, Wray DW, Hoff J, Carlier PG.** Human skeletal muscle intracellular oxygenation: the impact of ambient oxygen availability. *J Physiol* 571: 415–424, 2006.
  377. **Richardson RS, Noyszewski EA, Kendrick KF, Leigh JS, Wagner PD.** Myoglobin  $\text{O}_2$  desaturation during exercise Evidence of limited  $\text{O}_2$  transport. *J Clin Invest* 96: 1916–1926, 1995.
  378. **Rios E, Pizarro G.** Voltage sensor of excitation-contraction coupling in skeletal muscle. *Physiol Rev* 71: 849–908, 1991.
  379. **Roberts RA, Ghiasvand F, Parker D.** Biochemistry of exercise-induced metabolic acidosis. *Am J Physiol Regul Integr Comp Physiol* 287: R502–R516, 2004.
  380. **Romer LM, Haverkamp HC, Lovering AT, Pegelow DF, Dempsey JA.** Effect of exercise-induced arterial hypoxemia on quadriceps muscle fatigue in healthy humans. *Am J Physiol Regul Integr Comp Physiol* 290: R365–R375, 2006.

381. **Rose AJ, Hargreaves M.** Exercise increases  $\text{Ca}^{2+}$ -calmodulin-dependent protein kinase II activity in human skeletal muscle. *J Physiol* 553: 303–309, 2003.
382. **Rudolf R, Magalhaes PJ, Pozzan T.** Direct in vivo monitoring of sarcoplasmic reticulum  $\text{Ca}^{2+}$  and cytosolic cAMP dynamics in mouse skeletal muscle. *J Cell Biol* 173: 187–193, 2006.
383. **Rudolf R, Mongillo M, Magalhaes PJ, Pozzan T.** In vivo monitoring of  $\text{Ca}^{2+}$  uptake into mitochondria of mouse skeletal muscle during contraction. *J Cell Biol* 166: 527–536, 2004.
384. **Ruff RL.** Na current density at and away from end plates on rat fast- and slow-twitch skeletal muscle fibers. *Am J Physiol Cell Physiol* 262: C229–C234, 1992.
385. **Ruff RL.** Single-channel basis of slow inactivation of  $\text{Na}^{+}$  channels in rat skeletal muscle. *Am J Physiol Cell Physiol* 271: C971–C981, 1996.
386. **Ruff RL.** Sodium channel slow inactivation and the distribution of sodium channels on skeletal muscle fibres enable the performance properties of different skeletal muscle fibre types. *Acta Physiol Scand* 156: 159–168, 1996.
387. **Sacchetto R, Damiani E, Pallanca A, Margreth A.** Coordinate expression of  $\text{Ca}^{2+}$ -ATPase slow-twitch isoform and of beta calmodulin-dependent protein kinase in phospholamban-deficient sarcoplasmic reticulum of rabbit masseter muscle. *FEBS Lett* 481: 255–260, 2000.
388. **Sahlin K.** Metabolic factors in fatigue. *Sports Med* 13: 99–107, 1992.
389. **Sahlin K, Harris RC, Nylin B, Hultman E.** Lactate content and pH in muscle obtained after dynamic exercise. *Pflügers Arch* 367: 143–149, 1976.
390. **Sahlin K, Katz A, Broberg S.** Tricarboxylic acid cycle intermediates in human muscle during prolonged exercise. *Am J Physiol Cell Physiol* 259: C834–C841, 1990.
391. **Sahlin K, Ren JM.** Relationship of contraction capacity to metabolic changes during recovery from a fatiguing contraction. *J Appl Physiol* 67: 648–654, 1989.
392. **Saito A, Seiler S, Chu A, Fleischer S.** Preparation and morphology of sarcoplasmic reticulum terminal cisternae from rabbit skeletal muscle. *J Cell Biol* 99: 875–885, 1984.
393. **Salviati G, Sorenson MM, Eastwood AB.** Calcium accumulation by the sarcoplasmic reticulum in two populations of chemically skinned human muscle fibers Effects of calcium and cyclic AMP. *J Gen Physiol* 79: 603–632, 1982.
394. **Sandiford SD, Green HJ, Duhamel TA, Schertzer JD, Perco JD, Ouyang J.** Muscle Na-K-pump and fatigue responses to progressive exercise in normoxia and hypoxia. *Am J Physiol Regul Integr Comp Physiol* 289: R441–R449, 2005.
395. **Scherer NM, Deamer DW.** Oxidative stress impairs the function of sarcoplasmic reticulum by oxidation of sulfhydryl groups in the  $\text{Ca}^{2+}$ -ATPase. *Arch Biochem Biophys* 246: 589–601, 1986.
396. **Schertzer JD, Green HJ, Duhamel TA, Tupling AR.** Mechanisms underlying increases in SR  $\text{Ca}^{2+}$ -ATPase activity after exercise in rat skeletal muscle. *Am J Physiol Endocrinol Metab* 284: E597–E610, 2003.
397. **Schneider MF, Chandler WK.** Voltage dependent charge movement in skeletal muscle: a possible step in excitation-contraction coupling. *Nature* 242: 244–246, 1973.
398. **Schneider MF, Simon BJ.** Inactivation of calcium release from the sarcoplasmic reticulum in frog skeletal muscle. *J Physiol* 405: 727–745, 1988.
399. **Sejersted OM, Sjøgaard G.** Dynamics and consequences of potassium shifts in skeletal muscle and heart during exercise. *Physiol Rev* 80: 1411–1481, 2000.
400. **Shindoh C, Dimarco A, Thomas A, Manubay P, Supinski G.** Effect of N-acetylcysteine on diaphragm fatigue. *J Appl Physiol* 68: 2107–2113, 1990.
401. **Shkryl VM, Shirokova N.** Transfer and tunneling of  $\text{Ca}^{2+}$  from sarcoplasmic reticulum to mitochondria in skeletal muscle. *J Biol Chem* 281: 1547–1554, 2006.
402. **Shulman RG.** Glycogen turnover forms lactate during exercise. *Exercise Sport Sci Rev* 33: 157–162, 2005.
403. **Sjøgaard G, Adams RP, Saltin B.** Water and ion shifts in skeletal muscle of humans with intense dynamic knee extension. *Am J Physiol Regul Integr Comp Physiol* 248: R190–R196, 1985.
404. **Skurvydas A, Zachovajevs P.** Is post-tetanic potentiation, low frequency fatigue (LFF) and pre-contraction depression (PCD) co-existent in intermittent isometric exercises of maximal intensity? *Acta Physiol Scand* 164: 127–133, 1998.
405. **Smerdu V, Karsch-Mizrachi I, Campione M, Leinwand L, Schiaffino S.** Type IIx myosin heavy chain transcripts are expressed in type IIb fibers of human skeletal muscle. *Am J Physiol Cell Physiol* 267: C1723–C1728, 1994.
406. **Söderlund K, Greenhaff PL, Hultman E.** Energy metabolism in type I and type II human muscle fibres during short term electrical stimulation at different frequencies. *Acta Physiol Scand* 144: 15–22, 1992.
407. **Söderlund K, Hultman E.** ATP and phosphocreatine changes in single human muscle fibers after intense electrical stimulation. *Am J Physiol Endocrinol Metab* 261: E737–E741, 1991.
408. **Somlyo AV, Gonzalez-Serratos H, Shuman H, McCellan G, Somlyo AP.** Calcium release and ionic changes in the sarcoplasmic reticulum of tetanized muscle: an electron probe study. *J Cell Biol* 90: 577–594, 1981.
409. **Sostaric SM, Skinner SL, Brown MJ, Sangkabutra T, Medved I, Medley T, Selig SE, Fairweather I, Rutar D, McKenna MJ.** Alkalosis increases muscle  $\text{K}^{+}$  release, but lowers plasma  $[\text{K}^{+}]$  and delays fatigue during dynamic forearm exercise. *J Physiol* 570: 185–205, 2006.
410. **Spangenburg EE, Booth FW.** Molecular regulation of individual skeletal muscle fibre types. *Acta Physiol Scand* 178: 413–424, 2003.
411. **Spangenburg EE, Ward CW, Williams JH.** Effects of lactate on force production by mouse EDL muscle: implications for the development of fatigue. *Can J Physiol Pharmacol* 76: 642–648, 1998.
412. **Spriet LL, Lindinger MI, Heigenhauser GJ, Jones NL.** Effects of alkalosis on skeletal muscle metabolism and performance during exercise. *Am J Physiol Regul Integr Comp Physiol* 251: R833–R839, 1986.
413. **Spriet LL, Lindinger MI, Mckelvie RS, Heigenhauser GJ, Jones NL.** Muscle glycogenolysis and  $\text{H}^{+}$  concentration during maximal intermittent cycling. *J Appl Physiol* 66: 8–13, 1989.
414. **Spriet LL, Matsos CG, Peters SJ, Heigenhauser GJ, Jones NL.** Effects of acidosis on rat muscle metabolism and performance during heavy exercise. *Am J Physiol Cell Physiol* 248: C337–C347, 1985.
415. **Spriet LL, Söderlund K, Bergström M, Hultman E.** Skeletal muscle glycogenolysis, glycolysis, pH during electrical stimulation in men. *J Appl Physiol* 62: 616–621, 1987.
416. **Spriet LL, Watt MJ.** Regulatory mechanisms in the interaction between carbohydrate and lipid oxidation during exercise. *Acta Physiol Scand* 178: 443–452, 2003.
417. **Spruce AE, Standen NB, Stanfield PR.** Voltage-dependent ATP-sensitive potassium channels of skeletal muscle membrane. *Nature* 316: 736–738, 1985.
418. **St Pierre J, Buckingham JA, Roebuck SJ, Brand MD.** Topology of superoxide production from different sites in the mitochondrial electron transport chain. *J Biol Chem* 277: 44784–44790, 2002.
419. **St Pierre J, Drori S, Uldry M, Silvaggi JM, Rhee J, Jager S, Handschin C, Zheng K, Lin J, Yang W, Simon DK, Bachoo R, Spiegelman BM.** Suppression of reactive oxygen species and neurodegeneration by the PGC-1 transcriptional coactivators. *Cell* 127: 397–408, 2006.
420. **Stary CM, Hogan MC.** Impairment of  $\text{Ca}^{2+}$  release in single *Xenopus* fibers fatigued at various extracellular  $\text{Po}_2$ . *J Appl Physiol* 88: 1743–1748, 2000.
421. **Stary CM, Hogan MC.** Intracellular pH during sequential, fatiguing contractile periods in isolated single *Xenopus* skeletal muscle fibers. *J Appl Physiol* 99: 308–312, 2005.
422. **Steehgs K, Benders A, Oerlemans F, De Haan A, Heerschap A, Ruitenbeek W, Jost C, Van Deursen J, Perryman D, Pette D, Bruckwilder M, Koudijs J, Jap P, Veerkamp J, Wieringa B.** Altered  $\text{Ca}^{2+}$  responses in muscles with combined mitochondrial and cytosolic creatine kinase deficiencies. *Cell* 89: 93–103, 1997.
423. **Steele DS, Duke AM.** Metabolic factors contributing to altered  $\text{Ca}^{2+}$  regulation in skeletal muscle fatigue. *Acta Physiol Scand* 179: 39–48, 2003.
- 423a. **Stephenson DG.** Tubular system excitability: an essential component of excitation-contraction coupling in fast twitch fibres of



- vertebrate skeletal muscle. *J Muscle Res Cell Motil* 27: 259–274, 2006.
424. Stephenson DG, Lamb GD, Stephenson GM. Events of the excitation-contraction-relaxation (E-C-R) cycle in fast- and slow-twitch mammalian muscle fibres relevant to muscle fatigue. *Acta Physiol Scand* 162: 229–245, 1998.
  425. Stephenson DG, Nguyen LT, Stephenson GMM. Glycogen content and excitation-contraction coupling in mechanically skinned muscle fibres of the cane toad. *J Physiol* 519: 177–187, 1999.
  426. Stienen GJ, Van Graas IA, Elzinga G. Uptake and caffeine-induced release of calcium in fast muscle fibers of *Xenopus laevis*: effects of MgATP and  $P_i$ . *Am J Physiol Cell Physiol* 265: C650–C657, 1993.
  427. Stofan DA, Callahan LA, Dimarco AF, Nethery DE, Supinski GS. Modulation of release of reactive oxygen species by the contracting diaphragm. *Am J Respir Crit Care Med* 161: 891–898, 2000.
  428. Supinski G. Free radical induced respiratory muscle dysfunction. *Mol Cell Biochem* 179: 99–110, 1998.
  429. Sutton JR. Limitations to maximal oxygen uptake. *Sports Med* 13: 127–133, 1992.
  430. Sweeney HL, Bowman BF, Stull JT. Myosin light chain phosphorylation in vertebrate striated muscle: regulation and function. *Am J Physiol Cell Physiol* 264: C1085–C1095, 1993.
  431. Sweeney HL, Stull JT. Phosphorylation of myosin in permeabilized mammalian cardiac and skeletal muscle cells. *Am J Physiol Cell Physiol* 250: C657–C660, 1986.
  432. Tada M, Yamamoto T, Tonomura Y. Molecular mechanism of active calcium transport by sarcoplasmic reticulum. *Physiol Rev* 58: 1–79, 1978.
  433. Takagi Y, Shuman H, Goldman YE. Coupling between phosphate release and force generation in muscle actomyosin. *Philos Trans R Soc Lond B Biol Sci* 359: 1913–1920, 2004.
  434. Tallini YN, Ohkura M, Choi BR, Ji G, Imoto K, Doran R, Lee J, Plan P, Wilson J, Xin HB, Sanbe A, Gulick J, Mathai J, Robbins J, Salama G, Nakai J, Kotlikoff MI. Imaging cellular signals in the heart in vivo: Cardiac expression of the high-signal  $Ca^{2+}$  indicator GCaMP2. *Proc Natl Acad Sci USA* 103: 4753–4758, 2006.
  435. Tanabe T, Beam KG, Adams BA, Niidome T, Numa S. Regions of the skeletal muscle dihydropyridine receptor critical for excitation-contraction coupling. *Nature* 346: 567–569, 1990.
  436. Tavi P, Allen DG, Niemela P, Vuolteenaho O, Weckström M, Westerblad H. Calmodulin kinase modulates  $Ca^{2+}$  release in mouse skeletal muscle. *J Physiol* 551: 5–12, 2003.
  437. Terjung RL, Clarkson P, Eichner ER, Greenhaff PL, Hespel PJ, Israel RG, Kraemer WJ, Meyer RA, Spriet LL, Tarnopolsky MA, Wagenmakers AJ, Williams MH. American College of Sports Medicine roundtable. The physiological and health effects of oral creatine supplementation. *Med Sci Sports Exercise* 32: 706–717, 2000.
  438. Thompson LV, Balog EM, Riley DA, Fitts RH. Muscle fatigue in frog semitendinosus: alterations in contractile function. *Am J Physiol Cell Physiol* 262: C1500–C1506, 1992.
  439. Todd G, Butler JE, Taylor JL, Gandevia SC. Hyperthermia: a failure of the motor cortex and the muscle. *J Physiol* 563: 621–631, 2005.
  440. Trinh HH, Lamb GD. Matching of sarcoplasmic reticulum and contractile properties in rat fast- and slow-twitch muscle fibres. *Clin Exp Pharmacol Physiol* 33: 591–600, 2006.
  441. Tubman LA, Macintosh BR, Maki WA. Myosin light chain phosphorylation and posttetanic potentiation in fatigued skeletal muscle. *Pflügers Arch* 431: 882–887, 1996.
  442. Tupling R, Green H, Grant S, Burnett M, Ranney D. Postcontractile force depression in humans is associated with an impairment in SR  $Ca^{2+}$  pump function. *Am J Physiol Regul Integr Comp Physiol* 278: R87–R94, 2000.
  443. Turrens JF. Mitochondrial formation of reactive oxygen species. *J Physiol* 552: 335–344, 2003.
  444. Usher-Smith JA, Fraser JA, Bailey PS, Griffin JL, Huang CL. The influence of intracellular lactate and  $H^+$  on cell volume in amphibian skeletal muscle. *J Physiol* 573: 799–818, 2006.
  445. Van Beekvelt MC, Drost G, Rongen G, Stegeman DF, Van Engelen BG, Zwartz MJ.  $Na^+K^+$ -ATPase is not involved in the warming-up phenomenon in generalized myotonia. *Muscle Nerve* 33: 514–523, 2006.
  446. Van der Poel C, Edwards JN, Macdonald WA, Stephenson DG. Mitochondrial superoxide production in skeletal muscle fibers of the rat plus decreased fiber excitability. *Am J Physiol Cell Physiol* 292: C1353–C1360, 2007.
  447. Van der Poel C, Stephenson DG. Reversible changes in  $Ca^{2+}$ -activation properties of rat skeletal muscle exposed to elevated physiological temperatures. *J Physiol* 544: 765–776, 2002.
  448. Vandebrouck C, Duport G, Cognard C, Raymond G. Cationic channels in normal and dystrophic human myotubes. *Neuromuscular Disorders* 11: 72–79, 2001.
  449. Veech RL, Lawson JW, Cornell NW, Krebs HA. Cytosolic phosphorylation potential. *J Biol Chem* 254: 6538–6547, 1979.
  450. Verburb E, Dutka TL, Lamb GD. Long-lasting muscle fatigue: partial disruption of excitation-contraction coupling by elevated cytosolic  $Ca^{2+}$  concentration during contractions. *Am J Physiol Cell Physiol* 290: C1199–C1208, 2006.
  451. Verburb E, Murphy RM, Stephenson DG, Lamb GD. Disruption of excitation-contraction coupling and titin by endogenous  $Ca^{2+}$ -activated proteases in toad muscle fibres. *J Physiol* 564: 775–790, 2005.
  452. Vøllestad NK, Sejersted RB, Woods JJ, Bigland-Ritchie B. Motor drive and metabolic responses during repeated sub-maximal contractions in humans. *J Appl Physiol* 64: 1421–1427, 1988.
  453. Wallimann T, Wyss M, Brdiczka D, Nicolay K, Eppenberger HM. Intracellular compartmentation, structure and function of creatine kinase isoenzymes in tissues with high and fluctuating energy demands: the “phosphocreatine circuit” for cellular energy homeostasis. *Biochem J* 281: 21–40, 1992.
  454. Wallinga W, Meijer SL, Alberink MJ, Vliek M, Wienk ED, Ypey DL. Modelling action potentials and membrane currents of mammalian skeletal muscle fibres in coherence with potassium concentration changes in the T-tubular system. *Eur Biophys J* 28: 317–329, 1999.
  455. Walter G, Vandenborne K, Elliott M, Leigh JS. In vivo ATP synthesis rates in single human muscles during high intensity exercise. *J Physiol* 519: 901–910, 1999.
  456. Walton AG, Bodin WJ, Furedi H, Schwatz A. Nucleation of calcium phosphate from solution. *Can J Chem* 45: 2695–2701, 1967.
  457. Ward CW, Reiken S, Marks AR, Marty I, Vassort G, Lacampagne A. Defects in ryanodine receptor calcium release in skeletal muscle from post-myocardial infarct rats. *FASEB J* 17: 1517–1519, 2003.
  458. Watson PD, Garner RP, Ward DS. Water uptake in stimulated cat skeletal muscle. *Am J Physiol Regul Integr Comp Physiol* 264: R790–R796, 1993.
  459. Wegmann G, Zanolla E, Eppenberger HM, Wallimann T. In situ compartmentation of creatine kinase in intact sarcomeric muscle: the acto-myosin overlap zone as a molecular sieve. *J Muscle Res Cell Motil* 13: 420–435, 1992.
  460. Wehrens XH, Lehnart SE, Reiken S, Van Der NR, Morales R, Sun J, Cheng Z, Deng SX, De Windt LJ, Landry DW, Marks AR. Enhancing calstabin binding to ryanodine receptors improves cardiac and skeletal muscle function in heart failure. *Proc Natl Acad Sci USA* 102: 9607–9612, 2005.
  461. West W, Hicks A, Mckelvie R, O'Brien J. The relationship between plasma potassium, muscle membrane excitability and force following quadriceps fatigue. *Pflügers Arch* 432: 43–49, 1996.
  462. Westerblad H, Allen DG. Changes of myoplasmic calcium concentration during fatigue in single mouse muscle fibers. *J Gen Physiol* 98: 615–635, 1991.
  463. Westerblad H, Allen DG. Changes of intracellular pH due to repetitive stimulation of single fibres from mouse skeletal muscle. *J Physiol* 449: 49–71, 1992.
  464. Westerblad H, Allen DG. Myoplasmic free  $Mg^{2+}$  concentration during repetitive stimulation of single fibres from mouse skeletal muscle. *J Physiol* 453: 413–434, 1992.
  465. Westerblad H, Allen DG. The contribution of  $[Ca^{2+}]_i$  to the slowing of relaxation in fatigued single fibres from mouse skeletal muscle. *J Physiol* 468: 729–740, 1993.



466. **Westerblad H, Allen DG.** The influence of intracellular pH on contraction, relaxation and  $[Ca^{2+}]_i$  in intact single fibres from mouse muscle. *J Physiol* 466: 611–628, 1993.
467. **Westerblad H, Allen DG.** Relaxation,  $[Ca^{2+}]_i$  and  $[Mg^{2+}]_i$  during prolonged tetanic stimulation of intact, single fibers from mouse skeletal muscle. *J Physiol* 480: 31–43, 1994.
468. **Westerblad H, Allen DG.** The role of sarcoplasmic reticulum in relaxation of mouse muscle; effects of 2,5-di(tert-butyl)-1,4-benzohydroquinone. *J Physiol* 474: 291–301, 1994.
469. **Westerblad H, Allen DG.** The effects of intracellular injections of phosphate on intracellular calcium and force in single fibres of mouse skeletal muscle. *Pflügers Arch* 431: 964–970, 1996.
470. **Westerblad H, Allen DG, Lännergren J.** Muscle fatigue: lactic acid or inorganic phosphate the major cause? *News Physiol Sci* 17: 17–21, 2002.
471. **Westerblad H, Allen DG, Lee JA.** Measurements of intracellular calcium during fatiguing stimulation in single *Xenopus* muscle fibres. *Prog Clin Biol Res* 315: 231–232, 1989.
472. **Westerblad H, Bruton JD, Lännergren J.** The effect of intracellular pH on contractile function of intact, single fibres of mouse muscle declines with increasing temperature. *J Physiol* 500: 193–204, 1997.
473. **Westerblad H, Dahlstedt AJ, Lännergren J.** Mechanisms underlying reduced maximum shortening velocity during fatigue of intact, single fibres of mouse muscle. *J Physiol* 510: 269–277, 1998.
474. **Westerblad H, Duty S, Allen DG.** Intracellular calcium concentration during low-frequency fatigue in isolated single fibers of mouse skeletal muscle. *J Appl Physiol* 75: 382–388, 1993.
475. **Westerblad H, Lännergren J.** Force and membrane potential during and after fatiguing, intermittent tetanic stimulation of single *Xenopus* fibres. *Acta Physiol Scand* 128: 369–378, 1986.
476. **Westerblad H, Lännergren J.** Changes of the force-velocity relation, isometric tension and relaxation rate during fatigue in intact, single fibres of *Xenopus* skeletal muscle. *J Muscle Res Cell Motil* 15: 287–298, 1994.
477. **Westerblad H, Lännergren J.** Reduced maximum shortening velocity in the absence of phosphocreatine observed in intact fibres of *Xenopus* skeletal muscle. *J Physiol* 482: 383–390, 1995.
478. **Westerblad H, Lännergren J, Allen DG.** Slowed relaxation in fatigued skeletal muscle fibers of *Xenopus* and mouse. Contribution of  $[Ca^{2+}]_i$  and crossbridges. *J Gen Physiol* 109: 385–399, 1997.
479. **Westerblad H, Lee JA, Lamb AG, Bolsover SR, Allen DG.** Spatial gradients of intracellular calcium in skeletal muscle during fatigue. *Pflügers Arch* 415: 734–740, 1990.
480. **Westerblad H, Lee JA, Lännergren J, Allen DG.** Cellular mechanisms of fatigue in skeletal muscle. *Am J Physiol Cell Physiol* 261: C195–C209, 1991.
481. **Williams JH.** Contractile apparatus and sarcoplasmic reticulum function: effects of fatigue, recovery, elevated  $Ca^{2+}$ . *J Appl Physiol* 83: 444–450, 1997.
482. **Williams JH, Ward CW.** Reduced  $Ca^{2+}$ -induced  $Ca^{2+}$  release from skeletal muscle sarcoplasmic reticulum at low pH. *Can J Physiol Pharmacol* 70: 926–930, 1992.
483. **Williams JH, Ward CW, Klug GA.** Fatigue-induced alterations in  $Ca^{2+}$  and caffeine sensitivities of skinned muscle fibers. *J Appl Physiol* 75: 586–593, 1993.
484. **Williams JH, Ward CW, Spangenburg EE, Nelson RM.** Functional aspects of skeletal muscle contractile apparatus and sarcoplasmic reticulum after fatigue. *J Appl Physiol* 85: 619–626, 1998.
485. **Wilson GJ, Dos RC, Stephenson DG, Williams DA.** Effects of sulphydryl modification on skinned rat skeletal muscle fibres using 5,5'-dithiobis(2-nitrobenzoic acid). *J Physiol* 437: 409–430, 1991.
486. **Wilson JR.** Exercise intolerance in heart failure. Importance of skeletal muscle. *Circulation* 91: 559–561, 1995.
487. **Winegar BD, Haws CM, Lansman JB.** Subconductance block of single mechanosensitive ion channels in skeletal muscle fibers by aminoglycoside antibiotics. *J Gen Physiol* 107: 433–443, 1996.
488. **Wolosker H, Rocha JB, Engelder S, Panizzutti R, De Miranda J, De Meis L.** Sarco/endoplasmic reticulum  $Ca^{2+}$ -ATPase isoforms: diverse responses to acidosis. *Biochem J* 321: 545–550, 1997.
489. **Wredenberg A, Wibom R, Wilhelmsson H, Graff C, Wiener HH, Burden SJ, Oldfors A, Westerblad H, Larsson NG.** Increased mitochondrial mass in mitochondrial myopathy mice. *Proc Natl Acad Sci USA* 99: 15066–15071, 2002.
490. **Wright JR, McCloskey DI, Fitzpatrick RC.** Effects of muscle perfusion pressure on fatigue and systemic arterial pressure in human subjects. *J Appl Physiol* 86: 845–851, 1999.
491. **Wu KD, Lytton J.** Molecular cloning and quantification of sarcoplasmic reticulum  $Ca^{2+}$ -ATPase isoforms in rat muscles. *Am J Physiol Cell Physiol* 264: C333–C341, 1993.
492. **Xia R, Webb JA, Gnall LL, Cutler K, Abramson JJ.** Skeletal muscle sarcoplasmic reticulum contains a NADH-dependent oxidase that generates superoxide. *Am J Physiol Cell Physiol* 285: C215–C221, 2003.
493. **Zhang SJ, Bruton JD, Katz A, Westerblad H.** Limited oxygen diffusion accelerates fatigue development in mouse skeletal muscle. *J Physiol* 572: 551–559, 2006.
494. **Zhao X, Yoshida M, Brotto L, Takeshima H, Weisleder N, Hirata Y, Nosek TM, Ma J, Brotto M.** Enhanced resistance to fatigue and altered calcium handling properties of sarcalumenin knockout mice. *Physiol Gen* 23: 72–78, 2005.
495. **Zhi G, Ryder JW, Huang J, Ding P, Chen Y, Zhao Y, Kamm KE, Stull JT.** Myosin light chain kinase and myosin phosphorylation effect frequency-dependent potentiation of skeletal muscle contraction. *Proc Natl Acad Sci USA* 102: 17519–17524, 2005.
496. **Zuo L, Christofi FL, Wright VP, Bao S, Clanton TL.** Lipoygenase-dependent superoxide release in skeletal muscle. *J Appl Physiol* 97: 661–668, 2004.
497. **Zuo L, Christofi FL, Wright VP, Liu CY, Merola AJ, Berliner LJ, Clanton TL.** Intra- and extracellular measurement of reactive oxygen species produced during heat stress in diaphragm muscle. *Am J Physiol Cell Physiol* 279: C1058–C1066, 2000.
498. **Zuo L, Pasnicu S, Wright VP, Merola AJ, Clanton TL.** Sources for superoxide release: lessons from blockade of electron transport, NADPH oxidase, and anion channels in diaphragm. *Antioxid Redox Signal* 5: 667–675, 2003.